




Social vulnerability as a predictor of physical activity and screen time in European children

Isabel Iguacel^{1,2,3}  · Juan M. Fernández-Alvira^{1,4} · Karin Bammann^{5,6} · Charalambos Chadjiorgiou⁷ · Stefaan De Henauw⁸ · Regina Heidinger-Felső⁹ · Lauren Lissner¹⁰ · Nathalie Michels⁸ · Angie Page¹¹ · Lucia A. Reisch¹² · Paola Russo¹³ · Ole Sprengeler⁶ · Toomas Veidebaum¹⁴ · Claudia Börnhorst⁶ · Luis A. Moreno^{1,2,3,15} · On behalf of the IDEFICS consortium

Received: 2 March 2017 / Revised: 22 September 2017 / Accepted: 14 October 2017 / Published online: 24 October 2017
© Swiss School of Public Health (SSPH+) 2017

Abstract

Objectives To examine associations between social vulnerabilities and meeting physical activity (PA) and screen time (ST) recommendations during a 2-year follow-up.

Methods 13,891 children aged 2.0 to < 9.9 from eight European countries were assessed at baseline and 8482 children at follow-up. Children's sports club membership, PA and ST were collected via parental questionnaires. Moderate-to-vigorous physical activity (MVPA) was

objectively assessed with accelerometers. Performing at least 1 h of MVPA daily and spending less than 2 h of ST defined physically active and non-sedentary children, respectively. Vulnerable groups were defined at baseline as children whose parents had minimal social networks, from non-traditional families, with migrant origin or with unemployed parents. Logistic mixed-effects analyses were performed adjusting for classical socioeconomic indicators. **Results** Children whose parents had minimal social networks had a higher risk of non-compliance with PA recommendations (subjectively assessed) at baseline. Migrants and children with unemployed parents had longer ST. All vulnerable groups were less likely to be sports club members.

Claudia Börnhorst and Luis A. Moreno contributed equally and share the last authorship.

Electronic supplementary material The online version of this article (doi:[10.1007/s00038-017-1048-4](https://doi.org/10.1007/s00038-017-1048-4)) contains supplementary material, which is available to authorized users.

✉ Isabel Iguacel
iguacel@unizar.es

Juan M. Fernández-Alvira
juanmiguel.fernandez@cnic.es

Karin Bammann
bammann@uni-bremen.de

Charalambos Chadjiorgiou
chadjigeo@gmail.com

Stefaan De Henauw
Stefaan.DeHenauw@ugent.be

Regina Heidinger-Felső
felso.regina@pte.hu

Lauren Lissner
lauren.lissner@gu.se

Nathalie Michels
Nathalie.Michels@UGent.be

Angie Page
A.S.Page@bristol.ac.uk

Lucia A. Reisch
lr.ikl@cbs.dk

Paola Russo
prusso@isa.cnr.it

Ole Sprengeler
sprengel@leibniz-bips.de

Toomas Veidebaum
toomas.veidebaum@tai.ee

Claudia Börnhorst
boern@leibniz-bips.de

Luis A. Moreno
lmoreno@unizar.es

- 1 GENU (Growth, Exercise, NUtrition and Development) Research Group, Faculty of Health Sciences, University of Zaragoza, Edificio del SAI, C/Pedro Cerbuna s/n, 50009 Saragossa, Spain
- 2 Instituto Agroalimentario de Aragón (IA2), Saragossa, Spain
- 3 Instituto de Investigación Sanitaria Aragón (IIS Aragón), Saragossa, Spain

Conclusions Migrants and children with unemployed parents are at risk for excessive ST and all vulnerable groups have lower odds of being sports club members.

Keywords Vulnerable groups · Physical activity · Accelerometry · Screen time · Children · IDEFICS study

Abbreviations

PA	Physical activity
MVPA	Moderate to vigorous physical activity
IDEFICS	Identification and prevention of dietary- and lifestyle-induced health effects in children and infants
ST	Screen time
SES	Socio-economic status
T0	Baseline
T1	Follow-up after the intervention

Introduction

Regular physical activity (PA) during childhood is associated with improved musculoskeletal and cardiovascular health and lower adiposity (Janssen and Leblanc 2010; Strong et al. 2005). Insufficient PA and excessive screen time (ST) are independently associated with negative health outcomes (Ekelund et al. 2012). Therefore, increasing PA and decreasing sedentary time are public

health priorities. Current guidelines for children aged 5–18 recommend at least 1 h of moderate-to-vigorous physical activity (MVPA) per day (World Health Organization 2010) and to limit ST to no more than 2 h a day (American Academy of Pediatrics 2001). Despite these benefits, many children do not meet the recommended level of PA or ST (Konstabel et al. 2014).

Socio-economic status (SES) is an important determinant of health in adults but results for children and adolescents are less consistent (Drenowatz et al. 2010). Some studies showed that youth from higher SES are more physically active than youth from lower SES (Hanson and Chen 2007) while one study in China reported that high SES was positively associated with insufficient PA (Wang et al. 2016). One reason for these differences may be that associations may vary by domain of PA. The association between SES and sport may be different to that for active transport, both of which contribute to overall PA. Regarding ST results also seem ambiguous (Pate et al. 2011). Similarly, studies vary according to the sedentary measure used. Some studies showed children from high SES groups spent more time on non-screen sedentary behaviours (such as sitting or lying down) and those from low SES spent more time in screen-based sedentary behaviours (e.g. watching TV). However, no significant differences between children from low and high SES backgrounds were found for total sedentary time (sum of non-screen sedentary behaviours and ST) (Klitsie et al. 2013). Using subjective and objective methods, Foley et al. (2011) showed that, children and adolescents in New Zealand from areas of lower deprivation (i.e., higher SES) accumulated more total sedentary time than those from higher deprivation.

Inconsistent findings could be partially due to differential methods used to assess PA levels and sedentary time by e.g. subjective procedures such as questionnaires compared to objective measures, such as accelerometers (Raudsepp and Viira 2008). Accelerometers are more accurate at assessing total time spent engaging in PA at different intensity levels and recording inactive time (Hagströmer et al. 2010). However, questionnaires are preferable to assess domains of PA (e.g. transport, sport, leisure) and sedentary-related behaviours (Atkin et al. 2012).

The majority of studies to date have focused on the relationship between classical SES indicators (such as income, education and occupation), PA and sedentary behaviours (Tandon et al. 2012) but other indicators of social vulnerability, such as children whose parents lack a social network, children from non-traditional families (the child does not live with both parents), migrant children or children with unemployed parents, are rarely explored in the literature. Social vulnerabilities can be defined as social (e.g. migrant) and economic (e.g. unemployment)

⁴ Fundación Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC), Madrid, Spain

⁵ Institute for Public Health and Nursing Sciences (IPP), University of Bremen, Bremen, Germany

⁶ Leibniz Institute for Prevention Research and Epidemiology-BIPS, Bremen, Germany

⁷ Research and Education Foundation of Child Health, Strovolos, Cyprus

⁸ Department of Public Health, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium

⁹ Department of Paediatrics, University of Pécs, Pécs, Hungary

¹⁰ Section for Epidemiology and Social Medicine, University of Gothenburg, Gothenburg, Sweden, Gothenburg, Sweden

¹¹ Department of Exercise and Health Sciences, Centre for Sport, Exercise and Health, University of Bristol, Bristol, UK

¹² Department of Intercultural Communication and Management Copenhagen Business School, Copenhagen, Denmark

¹³ Institute of Food Sciences, National Research Council, Avellino, Italy

¹⁴ Department of Chronic Diseases, National Institute for Health Development, Tallinn, Estonia

¹⁵ Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición (CIBEROBn), Saragossa, Spain

situations that can increase the susceptibility to harm and that eventually amount to social disconnectedness (Haudenhuyse et al. 2012). These social vulnerable groups tend to adopt unhealthier behaviours and to be less active compared to non-vulnerable groups (Hawkins et al. 2009; Labree et al. 2014).

We hypothesized that children from vulnerable groups would have lower levels of PA and higher levels of ST compared to non-vulnerable groups due to financial constraints and negative experiences faced by vulnerable children. Some investigations reported that migrant children had lower levels of PA compared to native children as a result of the acculturation and a different body image perception (Labree et al. 2014). Non-traditional families could be at risk of being more inactive and of having lower sports participation levels because they might have lower modelling abilities and financial capacity compared to traditional families (Quarmby et al. 2011). Children with unemployed parents reported lower levels of PA and higher levels of ST compared to children with employed parents (Federico et al. 2009). Job loss raises TV-watching and since parents exert an impact on children, this may negatively affect children. Finally, we expect that children whose parents lack a social network could have a lower participation in PA and higher ST levels because of less access to resources and personal contact that could encourage activity levels.

To our knowledge no studies have examined a set of social vulnerabilities in the same population. Four vulnerable groups were investigated: children whose parents lack a social network; children from non-traditional families; migrant children and children with either one or both parents unemployed. This paper aims to explore (i) the cross-sectional and prospective associations between being a member (vs. non-member) of a vulnerable group at baseline and PA (reported and objectively assessed with accelerometers), sports club membership and ST, at two time points, in European children and (ii) the association of accumulated vulnerability (belonging to multiple vulnerable groups) with PA, sports club membership and ST at baseline. This will allow us to understand whether the disadvantages of socio-economic circumstances in European children are associated with unhealthy activity behaviours.

Methods

Design and study population

IDEFICS is a multi-centre prospective cohort study, including a school- and community-based obesity prevention intervention in eight European countries (Belgium,

Cyprus, Estonia, Germany, Hungary, Italy, Spain and Sweden). At baseline (T0), 16,228 children aged 2.0–9.9 years were examined from September 2007 to June 2008 (response rate 51%). The first follow-up (T1) took place 2 years later (September 2009–June 2010) when 11,038 children aged 4.0–11.9 years were re-examined. In all survey centres, recruitment was carried out at the community level. Parents of children eligible for inclusion were identified and recruited through local kindergartens and schools. The survey comprised anthropometrical measurements, examinations of children and parental self-completion questionnaires on lifestyle habits and dietary intakes of children. Standardised procedures were used by all survey centres. A detailed description is given by Ahrens et al. (2011).

Parents or legal guardians gave written informed consent for examinations and data collection for their children, while children expressed oral consent. Ethical approval was obtained from the research ethics authority of each participating centre.

Measurements

Physical activity and screen time assessed with a questionnaire

A parental questionnaire was used to collect a proxy measure of children's subjectively measured PA and ST (Burdette et al. 2004). Parents reported the total hours and minutes children spent playing outdoors during weekends and weekdays and the weekly duration their children spent doing sport in a sports club for a typical week in the previous month. Reported PA was calculated as: [(hours playing outdoors on weekdays * 5) + (hours playing outdoors on weekend days * 2) + weekly sports participation]/7. Thereafter, participants were classified depending on whether they met the current PA guidelines of < 1 h/day vs. \geq 1 h/day (World Health Organization 2010). Parents also reported children's sport club membership (dichotomized into belonging or not belonging to a sport club).

Moreover, parents reported hours of TV/DVD/video viewing and computer/games-console use for weekdays and weekend days by their children. Response options were: not at all; < 0.5; < 1; 1 to < 2; 2 to < 3; and \geq 3 h/day. Total ST per day was calculated as: (5 * weekday values + 2 * weekend values)/7. Participants were divided into two groups depending on whether they met current ST guidelines of \leq 2 h/day vs. > 2 h/day (American Academy of Pediatrics 2001).

Objectively measured MVPA

Children were instructed and asked to wear a uniaxial accelerometer (ActiGraph or ActiTrainer, ActiGraph, Pensacola, FL, USA) on a hip belt for at least 2 days including one weekend day and one-weekday (weekdays were weighted by five and weekend days by two and the sum was divided by seven). An average of 730 min of valid time was obtained in the final sample. To obtain comparable data despite differing valid times, adjusted MVPA was calculated by dividing raw minutes of MVPA by wear time and multiplying by 730 (Konstabel et al. 2014). Only children with a minimum duration of 8 h monitoring time per day were considered, where non-wear time was defined as at least 20 min of consecutive zeroes. The sampling epoch was set to 15 s but data were re-integrated into 60 s epochs for analysis. The duration of MVPA was determined according to the cut-offs of Evenson (Evenson et al. 2008).

Classical SES indicators as possible confounder

Education parents indicated the highest level of education of both themselves and their partners. The particular response categories for each country were coded according to the International Standard Classification of Education (UNESCO Institute for Statistics 1997) and re-categorized into: low (ISCED level 0–2), medium (ISCED level 3–4) and high (ISCED level 5–6) educational levels (UNESCO 1997).

Income parents provided information on the monthly net income of the household after taxes and deductions responding to nine country-specific categories (1: lowest income category to 9: highest income category). The category cut-offs were country-specific according to a scheme based on the median equivalent income, thus guaranteeing comparability between countries. The result was organised into three categories: low (1–3), medium (4–6) and high (7–9) income.

Occupation parents indicated their occupational position with 18 possible options, which were later transformed into the three-class version of the European Socioeconomic Classification (ESeC): working class, intermediate and salaried (Harrison and Rose 2006).

For occupation and education, the highest level of either the mother or the father was considered for the purpose of the study.

Vulnerable groups as predictors

Four vulnerabilities (dichotomised as vulnerable vs. non-vulnerable) were defined as our main exposures using baseline information from parent-reported questionnaires:

Social network based on the Single Item Measure of Social Support developed by Blake and McKay (1986) parents were asked how many persons they could rely on in case of need including their family. A minimal social network (vulnerable group) was assessed if the parental answer on the question was either ‘Nobody’ or ‘1 person’. Further answer categories were ‘2–3 persons’ and ‘more than 3 persons’ and were labelled as non-vulnerable (Bammann et al. 2013). This measure has been strongly associated with a composite social support index (Blake and McKay 1986).

Family structure If the child did not live with both his/her parents, the family was defined as a ‘non-traditional family’ (including single-parent families, stepparent families, living with grandparents or foster parents or in an institution).

Origin of parents A migrant background (vulnerable group) was assumed if one or both parents were born in a country different from where the study took place.

Employment status If at least one of the parents was unemployed or living on social assistance or welfare, the child was considered as belonging to the vulnerable group.

A total vulnerability score was calculated by adding up the numbers of vulnerabilities a child was exposed to. Six vulnerability indicators (minimal social network, non-traditional family, migrant, unemployed, low-income and low-education) were considered. Occupation status was not included as it was highly correlated with employment status. The vulnerability score ranged from 0 (the child had no vulnerabilities) to 6 (the child had all six vulnerability indicators) and was divided into four categories (three to six vulnerabilities, two vulnerabilities, one vulnerability and no vulnerability).

Weight categories

Anthropometric measurements were assessed at T0 according to standardised procedures in all participating countries. Barefoot body height was measured to the nearest 0.1 cm by trained staff using a portable stadiometer (SECA 225). Body weight in kg was measured by a child-adapted version of electronic scale TANITA BC 420 SMA with the children weighted in a fasting state and wearing only light clothes. Body mass index (BMI) was calculated by dividing body weight in kilograms by squared body height in metres and then transformed into an age- and gender-specific z-score (Cole et al. 1998). Weight groups were categorised using age and gender-specific cut points according to the criteria of the International Obesity Task Force (Cole and Lobstein 2012).

Sample size


Two analysis datasets were defined, one for the subjective and one for the objective measurements. Regarding the subjective measurements, 13,891 children were included for the cross-sectional analysis and 8482 children for the longitudinal analysis after excluding children with missing values in any of the outcomes (see Fig. 1). Children lost to follow-up belonged more often to the minimal social network group (12.0% vs. 9.0%), to non-traditional families (25% vs. 18.4%), migrants (16.5% vs. 12.8%) and unemployed parents (7.0% vs. 4.8%) than those included in the present study.

Concerning objective measurements, 9021 children had at least some valid accelerometer data at T0 but only 5892 children met the following quality requirements (Konstabel

et al. 2014): having at least 8 h daily wearing time for at least 2 days (1 weekend day and 1 weekday) using 60 s epoch. After 2 years of follow-up, only 2285 children measured at both T0 and T1 met the accelerometer quality criteria and were included in the longitudinal analysis (see Fig. 2). Children lost to follow-up belonged more often to non-traditional families (21.3% vs. 16.2%) and had more often a migrant background (16.5% vs. 12.8%) than those who were finally included in this study.


Statistical analyses

Logistic mixed-effects models were used to assess the cross-sectional and longitudinal associations between the four exposures (social network, family structure, migrant origin and employment status) and each outcome (meeting



Abbreviations: T0, baseline; T1, follow-up; PA, Physical Activity; ST, Screen Time.

Fig. 1 Selection of the study sample for subjective measurements (parent-reported questionnaire). T0 baseline, T1 follow-up, PA physical activity, ST screen time



Abbreviations: T0, baseline; T1, follow-up.

Fig. 2 Selection of the study sample for objective measurement (accelerometer). T0 baseline, T1 follow-up

recommendations for objectively and subjectively measured PA and ST, sports club membership). The reference category used was the healthiest behaviour for each outcome (subjective PA ≥ 1 h, ST ≤ 2 h, sports club membership and objective MVPA ≥ 1 h), respectively. The cross-sectional models were adjusted for baseline age, gender, BMI z-score and classical SES indicators; the objectively measured PA (MVPA) model was additionally adjusted for season. The longitudinal analyses were again adjusted for baseline age, gender, BMI z-score and classical SES indicators, but also for region (intervention versus control region) and baseline outcomes. A further analysis was conducted to estimate the accumulation of vulnerability at T0 and PA (subjectively and objectively assessed), sports club membership and ST. All models included a random kindergartens/school and a random country effect to account for the clustered study design.

Respondents with missing socioeconomic information may not be a random subset of population-based survey participants and excluding them may bias study results (Kim et al. 2007). Therefore, missing values of socioeconomic data were coded as a separate category.

Before model building, correlations among SES indicators were checked resulting in the exclusion of occupation status in models with employment status as main exposure to avoid collinearity problems.

The significance level was set at 0.01 to account at least partially for multiple testing. The analyses were performed using the Statistical Package for the Social Sciences (version 22.0; SPSS, Inc.).

Results

Table 1 summarises the distributions of predictors and background variables for the three parent reported outcomes (reported PA, ST and sports club membership) at T0 and T1 (see Table S1). Older children presented a higher percentage of meeting PA recommendations than younger children (88.6 and 86.1%, respectively), exceed ST recommendations (19.6 and 36.9%, respectively), and being a member of a sports club (27.1 and 58.5%, respectively). By sexes, males had a lower percentage of children reporting ≥ 1 h of PA (87%) and sports club membership (43.7%) than females (88 and 45%, respectively). By countries, Germany had the highest percentage of children being member of a sports club (58.3%) and Cyprus the lowest (38.5%).

Table 2 shows the distributions of predictors and background variables for the objectively measured PA (MVPA) at T0 and T1. The percentage of children reporting ≥ 1 h of MVPA was lower than subjectively measured PA.

Children from vulnerable groups and with missing values presented a lower percentage of meeting PA recommendations, a higher percentage of exceeding ST recommendations and a lower percentage as members of a sports club than non-vulnerable groups. Regarding T1, results were similar to T0 (see Table 1).

Table 3 and Table S2 present odds ratio (OR), 99% confidence interval (CI) and *p* values for the models assessing the cross-sectional and longitudinal associations between the four vulnerability indicators at T0 and the reported PA, ST and sports club membership at T0 and T1, respectively. Regarding T0, children whose parents had minimal social networks (OR = 1.30, [99% CI 1.10–1.61]) were more likely not to reach PA recommendations. Migrants (OR = 1.32, [99% CI 1.17–1.48]) and children with unemployed parents (OR = 1.33, [99% CI 1.07–1.66]) were less likely to meet ST recommendation. Those children whose parents had minimal social networks (OR 1.30, [99% CI 1.10–1.61]), non-traditional families (OR = 1.15, [99% CI 1.01–1.31]), migrants (OR = 1.49, [99% CI 1.33–1.68]) and children with unemployed parents (OR = 1.34, [99% CI 1.06–1.70]) were less likely to belong to a sports club. After 2-year follow-up, associations remained for non-traditional families and children with unemployed parents who were less likely to belong to a sports club at T1.

Table 4 shows the models assessing cross-sectional and longitudinal associations between the four vulnerability indicators at T0 and MVPA at T0 and T1, respectively. No associations were found between any of the social vulnerabilities and MVPA at T0 or T1.

Table S3 and S4 from supplementary material show the association between the accumulation of vulnerabilities and the four outcomes (reported and objectively assessed PA, ST and sports club membership) at T0. A higher number of vulnerabilities were not associated with a higher risk of non-compliance with PA recommendations (subjectively and objectively measured) but it was associated with a higher risk of non-compliance with ST recommendations, where the OR increased with the number of present vulnerabilities. Likewise, a greater number of vulnerabilities was associated with a lower likelihood of being a member of a sports club.

To estimate the change produced when including the classical SES indicators (full adjusted models), we added basic adjusted models (adjusted for baseline age, sex and BMI z-score) as supplementary material (see Table S5). ORs were greater when excluding classical SES compared to the full adjusted models. However, overall results remained unaltered.

Table 1 Description of the study population at T0, stratified by subjectively measured PA (divided into < 1 h of reported PA and ≥ 1 of reported PA), ST (divided into < 2 h of ST and ≥ 2 of ST) and belonging to a sport club (yes or no)

Total	T0						
	N (%)	Reported PA ^a		ST ^b		Sports club ^c	
		<1 h %	≥ 1 h %	>2 h %	≤ 2 h %	No %	Yes %
	13,891 (100%)						
Age groups (years)							
2 to < 6	6270 (45.1)	13.9	86.1	19.6	80.4	72.9	27.1
6 to < 10	7621 (54.9)	11.4	88.6	36.9	63.1	41.5	58.5
Gender of the child							
Male	7034 (50.6)	13.0	87.0	30.4	69.6	56.3	43.7
Female	6857 (49.4)	12.0	88.0	27.8	72.2	55.0	45.0
Country							
Italy	2062 (14.8)	20.0	80.0	33.1	66.9	59.4	40.6
Estonia	1548 (11.1)	12.7	87.3	44.3	55.7	50.0	50.0
Cyprus	1334 (9.6)	16.3	83.7	33.4	66.6	61.2	38.8
Belgium	1724 (12.4)	23.3	76.7	25.2	74.8	55.5	44.5
Sweden	1680 (12.1)	5.7	94.3	25.7	74.3	48.0	52.0
Germany	1790 (12.9)	9.8	90.2	22.5	77.5	41.7	58.3
Hungary	2377 (17.1)	5.9	94.1	28.6	71.4	72.1	27.9
Spain	1276 (9.9)	7.4	92.6	20.1	79.9	50.4	49.6
BMI categories							
Thinness	1576 (11.3)	13.0	87.0	23.8	76.2	64.5	35.5
Normal	9626 (69.3)	12.2	87.8	27.9	72.1	54.8	45.2
Overweight	1708 (12.3)	13.4	86.6	34.7	65.3	53.6	46.4
Obese	981 (7.1)	13.1	86.9	39.3	60.7	53.6	46.4
Social network ^d							
Missing	170 (1.2)	24.1	75.9	29.4	70.6	57.6	32.4
Minimal	1458 (10.5)	14.6	85.4	32.5	67.5	62.4	37.6
Strong	12,263 (88.3)	12.1	87.6	28.7	71.3	60.0	40.0
Family structure							
Missing	111 (8.0)	19.8	80.2	33.8	66.2	57.7	42.3
Non-traditional	2912 (21.0)	13.2	86.8	33.3	66.7	58.7	40.2
Traditional	10,868 (78.2)	12.3	87.7	27.8	72.2	54.5	45.5
Migrant status							
Missing	84 (0.6)	16.7	83.3	22.6	77.4	61.9	38.1
Migrant origin	1978 (14.2)	12.6	87.4	34.1	65.9	61.5	38.5
Native	11,829 (85.2)	12.5	87.5	28.3	71.4	54.6	45.4
Employment status							
Missing	182 (1.3)	15.9	84.1	36.3	63.7	66.5	33.5
Unemployed	782 (5.6)	13.2	86.8	35.4	64.6	67.0	33.0
Employed	12,927 (93.1)	12.5	87.5	28.6	71.4	54.8	45.2
Parental education							
Missing	96 (0.7)	20.8	79.2	32.3	67.7	66.7	33.3
Low	1112 (8.0)	12.2	87.8	39.7	60.3	70.9	29.1
Medium	6964 (50.1)	11.8	88.2	33.0	67.0	57.0	43.0
High	5719 (41.2)	13.4	86.6	22.1	77.9	50.9	49.1
Income							
Missing	921 (6.6)	17.4	83.3	22.6	77.4	61.9	38.1
Low	4743 (34.1)	12.6	87.4	34.1	65.9	61.5	38.5
Medium	3704 (26.7)	12.5	87.5	28.3	71.4	54.6	45.4
High	4523 (32.6)	11.0	89.0	25.8	74.2	50.0	50.0

Table 1 continued

Total	T0						
	N (%)	Reported PA ^a		ST ^b		Sports club ^c	
		<1 h %	≥ 1 h %	>2 h %	≤ 2 h %	No %	Yes %
	13,891 (100%)						
Parental occupation							
Missing	485 (3.5)	15.1	84.9	34.8	65.2	67.4	32.6
Working class	4603 (33.1)	12.0	88.0	35.2	64.8	65.3	34.7
Intermediate	4999 (36.0)	13.0	87.0	26.1	73.9	52.2	47.8
Salariat	3804 (27.4)	12.3	87.7	24.9	75.1	47.1	52.9

Study population: children from 8 European countries aged 2.0–9.9 years examined from September 2009–June 2010

T0 baseline, PA physical activity, ST screen time, h hour(s)

^aReported PA: sum of hours that children spent playing outdoors (weekdays and weekend days) and weekly participation in sport club activities

^bScreen time: total number of hours usually spent watching TV, videos or DVD and playing on the computer or games console

^cSport club membership

^dSocial network was assessed with the question how many persons they could rely on in case of need including their family: minimal (0–1 person) and strong (> 2 persons)

Discussion

This paper investigated the association between PA (objectively and subjectively assessed), sports club membership, ST and social vulnerabilities over a 2-year period in children aged 2.0–9.9 years participating in a large European cohort study. Vulnerable children presented a higher risk of showing excessive ST cross-sectionally and tended to be less active at sports clubs cross-sectionally and longitudinally, compared to non-vulnerable groups. Regarding PA, our results did not show a strong association with social vulnerability indicators. Only those children whose parents reported to have minimal social networks were found to be at higher risk of non-compliance with subjectively assessed PA recommendations.

Adjusting for classical SES indicators allowed investigation of whether the associations between social vulnerabilities and ST/PA were independent of classical SES indicators or whether only the classical SES indicators were finally relevant in the model. We observed that associations may be partly explained by classical SES variables but still independent of classical SES indicators. Therefore, belonging to a vulnerable group seems to be an independent factor of excessive ST and lower participation/activity at sports clubs. A greater effect of the vulnerabilities was observed in cross-sectional analyses as opposed to longitudinal analyses. Consequently, current vulnerability (at the time of outcome assessment) seems the most relevant one for children's PA and ST.

The findings of our study are in line with previous research (McMillan et al. 2015; Singhammer et al. 2015) despite some differences.

Regarding family structure, no significant associations were found between children from non-traditional families and PA or ST, which is in agreement with some studies (McMillan et al. 2015; Singhammer et al. 2015). However, other studies have reported that children from non-traditional families accumulate more ST and a higher risk of not meeting PA recommendations as a result of differences in role modelling abilities and financial capacity (Bagley et al. 2006; Quarmby et al. 2011).

Concerning migrant status, we found statistical differences between migrant children and exceeding ST recommendations at baseline. The acculturation in the host society acquiring Western lifestyle characterized by lower levels of PA and higher levels of sedentary behaviours and different body image perceptions maintained from the country of origin could be the reason of differences found between migrant and native children. However, no association was found between migrant children and not meeting PA recommendations. Similar to our study, Puder et al. (2013) showed that migrant children had a significantly higher amount of ST compared with children born in the country of measurement. Contrary to what we observed, it was showed that PA levels in children were significantly lower among migrant children compared to children in the native population (Labree et al. 2014).

Children whose parents were unemployed were more likely to exceed ST recommendations at baseline compared

Table 2 Description of the study population, stratified by objectively measured PA (MVPA) at T0 and T1

Total	MVPA T0		MVPA T1			
	N (%)	<1 h %	≥1 h %	N (%)	<1 h %	≥1 h %
Age groups (years)						
2 to < 6	2471 (41.9)	87.6	12.4	993 (43.5)	75.6	24.4
6 to < 12	3421 (58.1)	77.9	22.1	1292 (56.5)	82.0	18.0
Country						
Italy	855 (14.5)	93.5	6.5	335 (14.7)	92.8	7.2
Estonia	931 (15.8)	81.5	18.5	565 (24.7)	81.2	18.8
Cyprus	402 (6.8)	92.0	8.0	83 (3.6)	94.0	6.0
Belgium	432 (7.3)	86.6	14.4	188 (8.2)	85.6	14.4
Sweden	486 (8.2)	73.5	26.5	61 (2.7)	80.3	19.7
Germany	913 (15.5)	71.7	28.3	177 (7.7)	74.0	26.0
Hungary	720 (12.2)	82.5	17.5	90 (3.9)	86.7	13.3
Spain	1153 (10.6)	80.3	19.7	786 (34.4)	69.2	30.8
BMI categories						
Thinness	597 (10.1)	85.1	14.9	226 (9.9)	75.7	24.3
Normal	4074 (69.1)	79.7	20.3	1603 (70.2)	81.3	18.7
Overweight	779 (13.2)	85.6	14.4	292 (12.8)	78.8	21.2
Obese	442 (7.5)	91.9	8.1	164 (7.2)	92.1	7.9
Social network ^a						
Missing	277 (4.7)	87.4	12.6	61 (2.6)	83.6	16.4
Minimal	640 (10.9)	82.0	18.0	278 (12.2)	81.3	18.7
Strong	4975 (84.4)	81.7	18.3	1946 (86.2)	78.8	21.2
Family structure						
Missing	247 (4.2)	89.1	10.9	50 (2.2)	88.0	12.0
Non-traditional family	1138 (19.3)	82.9	17.1	370 (16.2)	80.3	19.7
Traditional family	4507 (76.5)	81.4	18.6	1865 (81.6)	78.8	21.2
Migrant status						
Missing	126 (2.1)	82.5	17.5	34 (1.5)	82.4	17.6
Migrant origin	792 (13.4)	82.8	17.2	237 (10.4)	84.8	15.2
Native	4974 (84.4)	81.8	18.2	2014 (88.1)	78.6	21.4
Employment status						
Missing	290 (4.9)	88.3	11.7	66 (2.9)	89.4	10.6
Unemployed	338 (5.7)	80.5	19.5	123 (5.4)	79.7	20.3
Employed	5264 (89.3)	81.7	18.3	2096 (91.7)	78.9	21.1
Parental education						
Missing	135 (2.3)	84.4	15.6	35 (1.5)	82.9	17.1
Low	507 (8.6)	82.6	17.4	146 (6.4)	80.8	19.2
Medium	3088 (52.4)	84.4	15.6	1263 (55.3)	80.0	20.0
High	2162 (36.7)	82.4	17.6	841 (36.8)	77.6	22.4
Income						
Missing	588 (10.0)	85.0	15.0	169 (7.4)	85.2	14.8
Low	1823 (30.9)	83.9	16.1	637 (27.9)	84.6	15.4
Medium	1550 (26.3)	79.7	20.3	625 (27.4)	75.0	25.0
High	2162 (36.6)	81.0	19.0	854 (37.4)	77.2	22.8
Parental occupation						
Missing	404 (6.9)	86.1	13.9	85 (3.7)	84.7	15.3
Working class	1881 (31.9)	82.3	17.7	683 (29.9)	81.7	18.3
Intermediate	2056 (34.9)	81.8	18.2	844 (36.9)		22.9

Table 2 continued

Total	MVPA T0			MVPA T1		
	N (%)	<1 h %	≥1 h %	N (%)	<1 h %	≥1 h %
	5892 (100%)			2285 (100%)		
Salariat	1551 (26.3)	80.8	19.2	673 (29.5)	78.8	21.2

Study population: children from 8 European countries aged 2.0–9.9 years examined at T0 from September 2007 to June 2008 and re-examined 2 years later (T1) from September 2009 to June 2010 when children aged 4.0–11.9 years

T0 baseline, T1 follow-up, MVPA moderate to vigorous physical activity (objectively measured), ST screen time, h hour(s)

^aSocial network was assessed with the question how many persons they could rely on in case of need including their family: minimal (0–1 person) and strong (> 2 persons)

Table 3 Cross-sectional and longitudinal associations between social vulnerability indicators and the three reported outcomes (subjectively measured via questionnaires) for the adjusted models

	Outcome at T0 ^{a,b}								
	Reported PA ^d			ST ^e			Sports club member ^f		
	OR	99% CI	<i>p</i> value	OR	99% CI	<i>p</i> value	OR	99% CI	<i>p</i> value
Social network ^g									
Missing	1.62	1.11–2.35	0.012	0.92	0.64–1.31	0.657	1.51	0.94–2.43	0.024
Minimal	1.30	1.10–1.61	0.002	1.05	0.92–1.19	0.436	1.23	1.09–1.47	0.001
Strong	1.00			1.00			1.00		
Family structure									
Missing	1.47	0.77–2.80	0.617	1.11	0.64–1.93	0.617	1.00	0.57–1.74	0.990
Non-traditional	0.97	0.84–1.19	0.952	1.00	0.88–1.13	0.921	1.15	1.01–1.31	0.007
Traditional	1.00			1.00			1.00		
Migrant status									
Missing	0.59	0.26–1.33	0.203	0.42	0.21–1.08	0.018	0.79	0.41–1.51	0.485
Migrant origin	1.00	0.85–1.17	0.971	1.32	1.17–1.48	< 0.001	1.49	1.33–1.68	< 0.001
Native	1.00			1.00			1.00		
Employment status									
Missing	1.12	0.58–2.17	0.663	1.15	0.79–1.86	0.456	1.00	0.60–1.65	0.950
Unemployed	1.23	0.97–1.66	0.076	1.33	1.07–1.66	0.001	1.34	1.06–1.70	0.001
Employed	1.00			1.00			1.00		

Results from the logistic mixed-effects models: odds ratios (OR), 99% confidence intervals (CI) and *p* values are shown. Study population: children from eight European countries aged 2.0–9.9 years examined at T0 from September 2007 to June 2008 and re-examined 2 years later (T1) from September 2009 to June 2010 when children aged 4.0–11.9 years

Bold indicates statistical significance

T0 baseline, T1 follow-up, PA physical activity, ST screen time

^aAll models at T0 were adjusted for baseline age, gender, baseline classical SES indicators (education, income and occupation) and z-score of BMI by (Cole and Lobstein 2012)

^bAll models include random effects (school, country) to account for the study design

^cModels at T1 were additionally adjusted for study region (intervention v. control) and baseline outcomes (reported PA, ST and sport club membership at T0, respectively)

^dReported PA: sum of hours that children spent playing outdoors (weekdays and weekend days) and weekly participation in sport club activities. Reference: Reported PA ≥ 1 h

^eScreen time: total number of hours usually spent watching TV, videos or DVD and playing on the computer or games console. Reference: ST ≤ 2 h

^fSport club membership. Reference: yes

^gSocial network was assessed with the question how many persons they could rely on in case of need including their family: minimal (0–1 person) and strong (> 2 persons)

Table 4 Cross-sectional and longitudinal associations between vulnerabilities indicators and MVPA (objectively measured with accelerometers) at baseline and follow up for the adjusted models

	MVPA T0 ^{a,b}			MVPA T1 ^{b,c}		
	OR	99% CI	<i>p</i> value	OR	99% CI	<i>p</i> value
Social network ^d						
Missing	1.14	0.57–2.27	0.615	0.78	0.31–1.97	0.603
Minimal	0.90	0.66–1.23	0.401	1.30	0.92–1.83	0.133
Strong	1.00			1.00		
Family structure						
Missing	1.44	0.68–3.07	0.210	0.87	0.59–1.71	0.499
Non-traditional family	1.05	0.81–1.36	0.606	0.82	0.53–1.27	0.244
Traditional family	1.00			1.00		
Migrant status						
Missing	0.40	0.09–1.80	0.119	0.70	0.03–1.21	0.783
Migrant origin	1.17	0.86–1.58	0.181	1.60	0.91–2.72	0.033
Native	1.00			1.00		
Employment status						
Missing	1.35	0.62–2.96	0.322	0.76	0.32–2.28	0.156
Unemployed	0.99	0.66–1.49	0.970	1.08	0.56–2.10	0.759
Employed	1.00			1.00		

Results from the logistic mixed-effects models: *p* values, odds ratios (OR) and 99% confidence intervals (CI) are shown

Study population: children from 8 European countries aged 2–9.9 years examined at T0 from September 2007 to June 2008 and re-examined 2 years later (T1) from September 2009 to June 2010 when children aged 4–11.9 years

T0 baseline, T1 follow-up, MVPA moderate-to-vigorous physical activity

^aModels at T0 were adjusted for baseline age, gender, season, baseline classical SES indicators (education, income and occupation) and z-score of BMI by (Cole and Lobstein 2012)

^bAll models include random effects (school, country) to account for the study design

^cModels at T1 were additionally adjusted for study region (intervention vs. control) and MVPA at baseline

^dSocial network was assessed with the question how many persons they could rely on in case of need including their family: minimal (0–1 person) and strong (> 2 persons)

to non-unemployed parents. These conclusions were confirmed by previous papers (Hawkins et al. 2009; van Rossem et al. 2012). Unemployed people are at a higher risk of depression and inactivity compared to employed people. Since parents are important role models for children this could lead to lower activity levels in children (Van Domelen et al. 2011). Nonetheless, our results did not show any association between children with unemployed parents and being at higher risk of not meeting PA recommendations, like other investigations have demonstrated (Federico et al. 2009).

Children whose parents had minimal social networks had a higher risk of non-compliance with PA recommendations (subjectively assessed) at baseline but they did not show a higher risk of exceeding ST recommendations. Not only parents but their networks can influence children's behaviours. Therefore, children whose parents have large social networks could have a positive influence for

performing higher levels of PA. To our knowledge, no studies have investigated the associations between parent's social network and children's PA and ST.

In line with previous studies, we found that all vulnerable groups were less likely to participate at sports clubs than children from non-vulnerable groups at baseline and follow-up (McMillan et al. 2016; Toftegaard-Støckel et al. 2010). These associations were rather weak for children from non-traditional families and higher for children with unemployed parents.

Some limitations of the present study should be acknowledged. First, the IDEFICS study is not representative of the European population or of the participating countries. Each survey centre only covered a delimited geographic area within a country making extrapolation of the results difficult and only a sub-sample of the participants wore an accelerometer. Furthermore, a selection bias cannot be precluded as the children lost to follow-up had

more social vulnerabilities at baseline and as voluntary participation might be less frequent from very high or very low SES families. Besides, since self-reported PA usually overestimates total PA compared to accelerometers, subjective PA data should be interpreted with caution. It is questionable how reliably the duration of outdoor-play and sports club membership capture total PA and how reliable the dichotomization of meeting the PA guidelines is according to self-reported PA. On the other side, accelerometers may underestimate the overall activity because they cannot accurately capture activities that are not step-based (such as swimming or cycling) (Colley et al. 2011). Therefore, MVPA may be diminished, which may partly explains the current results as associations would be attenuated. Moreover, valid data on accelerometers was considered when children had at least 2 days of recording time (including one weekend day and one-week day) with a minimum 8-h duration of monitoring time per day, which could be insufficient for a correct assessment of whether they meet the PA guidelines. Finally, even though we have controlled for several potential confounders, we cannot preclude unmeasured confounding e.g. through parents' health status, parents' mental health and other socio-cultural factors.

A particular strength of this study is that to our knowledge, no research has been done concerning the association of vulnerabilities such as social network, family structure and unemployment status with objectively and subjectively assessed PA and ST in children in a longitudinal study. Having two measures of PA (subjectively and objectively assessed) provide different information. For example, sports club participation usually requires regular payments and it has hence other barriers than playing on a playground. Accelerometers could register both activities but it could not distinguish these differences. The large sample size of eight countries following standardised procedures is also a strength.

Future studies may investigate children with a different country of origin and family structure in more depth to help identify children at higher risk of low PA and high ST. Moreover, more studies including both subjective and objective measures of PA levels and sedentary behaviours are needed to test different constructs which provide additional information and compare possible discrepancies in results to analyse the causes.

Conclusion

The results suggest a higher risk for excessive ST cross-sectionally in children with unemployed parents and migrants as well as lower odds of being a member in a sports club cross-sectionally and longitudinally in all vulnerable groups independent of family income, parental

occupation and parental education. However, no associations were found between any of the social vulnerabilities and objectively assessed PA. Policy makers should focus on decreasing ST sedentary behaviours among vulnerable groups as well as on offering subsidised access to external exercise, fitness, sports clubs and facilities.

Acknowledgements This work has been done as part of the IDEFICS Study (<http://www.idefics.eu>). The authors gratefully acknowledge the financial support of the European Community within the Sixth RTD Framework Programme Contract No. 016181 (FOOD). I. I has been financed by the FPU (Grant Reference FPU014/00922) Pre-doctoral Programs (Spanish Ministry of Education and Science). We thank the IDEFICS children and their parents who generously volunteered and participated in this project.

Authors' contributions The authors' contributions were as follows: II carried out the statistical analysis and drafted the manuscript. CB designed the statistical analyses. KB, JMF-A, NM, LL, and OS, developed the measurement instruments; SDH, RF, LAM, AP, LAR, PR, and TV supervised the national data collection procedures. All authors read and critically reviewed the manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that there are no conflicts of interest regarding this manuscript.

References

- Ahrens W, Bammann K, Siani A et al (2011) The IDEFICS cohort: design, characteristics and participation in the baseline survey. *Int J Obes* 35(Suppl 1):S3–15. doi:10.1038/ijo.2011.30
- American Academy of Pediatrics (2001) Children, adolescents, and television. *Pediatrics* 107:423–426. doi:10.1542/peds.107.2.423
- Atkin AJ, Gorely T, Clemen SA et al (2012) Methods of measurement in epidemiology: sedentary behaviour. *Int J Epidemiol* 41:1460–1471. doi:10.1093/ije/dys118
- Bagley S, Salmon J, Crawford D (2006) Family structure and children's television viewing and physical activity. *Med Sci Sports Exerc* 38:910–918. doi:10.1249/01.mss.0000218132.68268.f4
- Bammann K, Gwozdz W, Lanfer A et al (2013) Socioeconomic factors and childhood overweight in Europe: results from the multi-centre IDEFICS study. *Pediatr Obes* 8:1–12. doi:10.1111/j.2047-6310.2012.00075.x
- Blake RL Jr, McKay DA (1986) A single-item measure of social supports as a predictor of morbidity. *J Fam Pract* 22:82–84
- Burdette HL, Whitaker RC, Daniels SR (2004) Parental report of outdoor playtime as a measure of physical activity in preschool-aged children. *Arch Pediatr Adolesc Med* 158:353–357. doi:10.1001/archpedi.158.4.353
- Cole TJ, Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 7:284–294. doi:10.1111/j.2047-6310.2012.00064.x
- Cole TJ, Freeman JV, Preece MA (1998) British 1990 growth reference centiles for weight, height, body mass index and head circumference fitted by maximum penalized likelihood-. *Stat Med* 17:407–429
- Colley RC, Garriguet D, Janssen I, Craig CL, Clarke J, Tremblay MS (2011) Physical activity of Canadian adults: accelerometer

- results from the 2007 to 2009 Canadian Health Measures Survey. *Health Rep* 22:7–14
- Drenowatz C, Eisenmann JC, Pfeiffer KA, Welk G, Heelan K, Gentile D, Walsh D (2010) Influence of socio-economic status on habitual physical activity and sedentary behavior in 8- to 11-year old children. *BMC Public Health* 10:214. doi:10.1186/1471-2458-10-214
- Ekelund U, Luan J, Sherar LB, Esliger DW, Griew P, Cooper A (2012) Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA* 307:704–712. doi:10.1001/jama.2012.156
- Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG (2008) Calibration of two objective measures of physical activity for children. *J Sports Sci* 26:1557–1565. doi:10.1080/02640410802334196
- Federico B, Falese L, Capelli G (2009) Socio-economic inequalities in physical activity practice among Italian children and adolescents: a cross-sectional study. *Z. Gesundh Wiss* 17:377–384. doi:10.1007/s10389-009-0267-4
- Foley LS, Maddison R, Jiang Y, Olds T, Ridley K (2011) It's not just the television: survey analysis of sedentary behaviour in New Zealand young people. *Int J Behav Nutr Phys Act* 8:132. doi:10.1186/1479-5868-8-132
- Hagströmer M, Ainsworth BE, Oja P, Sjöström M (2010) Comparison of a subjective and an objective measure of physical activity in a population sample. *J Phys Act Health* 7:541–550
- Hanson MD, Chen E (2007) Socioeconomic status and health behaviors in adolescence: a review of the literature. *J Behav Med* 30:263–285. doi:10.1007/s10865-007-9098-3
- Harrison E, Rose D (2006) The european socio-economic classification (ESeC) user guide. University of Essex, Colchester. <https://www.iser.essex.ac.uk/files/esecc/guide/docs/UserGuide.pdf>. Accessed 20 July 2016
- Haudenhuyse RP, Theeboom M, Coalter F (2012) The potential of sports-based social interventions for vulnerable youth: implications for sport coaches and youth workers. *J Youth Stud* 15(4):437–454. doi:10.1080/13676261.2012.663895
- Hawkins SS, Cole TJ, Law C (2009) Examining the relationship between maternal employment and health behaviours in 5-year-old British children. *J Epidemiol Community Health* 63:999–1004. doi:10.1136/jech.2008.084590
- Janssen I, Leblanc AG (2010) Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act* 7:40. doi:10.1186/1479-5868-7-40
- Kim S, Egarter S, Cubbin C, Takahashi ER, Braveman P (2007) Potential implications of missing income data in population-based surveys: an example from a postpartum survey in California. *Public Health Rep* 122:753–763
- Klitsie T, Corder K, Visscher TL, Atkin AJ, Jones AP, van Sluijs EM (2013) Children's sedentary behaviour: descriptive epidemiology and associations with objectively-measured sedentary time. *BMC Public Health* 13:1092. doi:10.1186/1471-2458-13-1092
- Konstabel K, Veidebaum T, Verbestel V et al (2014) Objectively measured physical activity in European children: the IDEFICS study. *Int J Obes* 38(Suppl 2):S135–S143. doi:10.1038/ijo.2014.144
- Labree W et al (2014) Physical activity differences between children from migrant and native origin. *BMC Public Health* 14:819. doi:10.1186/1471-2458-14-819
- McMillan R, McIsaac M, Janssen I (2015) Family structure as a predictor of screen time among youth. *PeerJ* 3:e1048. doi:10.7717/peerj.1048
- McMillan R, McIsaac M, Janssen I (2016) Family structure as a correlate of organized sport participation among youth. *PLoS ONE* 11:e0147403. doi:10.1371/journal.pone.0147403
- Pate RR, Mitchell JA, Byun W, Dowda M (2011) Sedentary behaviour in youth. *Br J Sports Med* 45:906–913. doi:10.1136/bjsports-2011-090192
- Puder J, Pinto AM, Bonvin A, Bodenman P, Munsch S, Kriemler S, Marques-Vidal P (2013) Health-related quality of life in migrant preschool children. *BMC Public Health* 13:384. doi:10.1186/1471-2458-13-384
- Quarmby T, Dagkas S, Bridge M (2011) Associations between children's physical activities, sedentary behaviours and family structure: a sequential mixed methods approach. *Health Educ Res* 26:63–76. doi:10.1093/her/cyq071
- Raudsepp L, Viira R (2008) Changes in physical activity in adolescent girls: a latent growth modelling approach. *Acta Paediatr* 97:647–652. doi:10.1111/j.1651-2227.2008.00748.x
- Singhammer J, Ried-Larsen M, Møller NC, Lund-Kristensen P, Froberg K, Andersen LB (2015) Single parent status and children's objectively measured level of physical activity. *Sports Med Open* 1:10. doi:10.1186/s40798-015-0020-1
- Strong WB, Malina RM, Blimkie CJ et al (2005) Evidence based physical activity for school-age youth. *J Pediatr* 146:732–737. doi:10.1016/j.jpeds.2005.01.055
- Tandon PS, Zhou C, Sallis JF, Cain KL, Frank LD, Saelens BE (2012) Home environment relationships with children's physical activity, sedentary time, and screen time by socioeconomic status. *Int J Behav Nutr Phys Act* 9:88. doi:10.1186/1479-5868-9-88
- Toftegaard-Stöckel J, Strandbu A, Solenes O, Jørgensen P, Fransson K (2010) Sport for children and youth in the Scandinavian countries. *Sport Soc* 13:625–642. doi:10.1080/17430431003616332
- UNESCO Institute for Statistics (1997) International Standard Classification of Education (ISCED). UNESCO, Montreal. 2006. http://www.unesco.org/education/information/nfsunesco/doc/isced_1997.htm. Accessed 15 July 2016
- Van Domelen DR, Koster A, Caserotti P et al (2011) Employment and physical activity in the U.S. *Am J Prev Med* 41:136–145. doi:10.1016/j.amepre.2011.03.019
- van Rossem L, Vogel I, Moll HA, Jaddoe VW, Hofman A, Mackenbach JP, Raat H (2012) An observational study on socio-economic and ethnic differences in indicators of sedentary behavior and physical activity in preschool children. *Prev Med* 54:55–60. doi:10.1016/j.yjmed.2011.10.016
- Wang X, Hui Z, Terry PD et al (2016) Correlates of insufficient physical activity among junior high school students: a cross-sectional study in Xi'an, China. *Int J Environ Res Public Health* 13:397. doi:10.3390/ijerph13040397
- World Health Organization (2010) Global recommendations on Physical activity for Health. World Health Organization, Geneva. http://apps.who.int/iris/bitstream/10665/44399/1/9789241599979_eng.pdf. Accessed 26 June 2016