



OPEN ACCESS

EDITED BY
 Jens Bucksch,
 Heidelberg University of Education,
 Germany

REVIEWED BY
 Dagmar Sigmundová,
 Palacký University, Olomouc, Czechia
 Roshelle Payes,
 Nemours Foundation, United States

*CORRESPONDENCE
 Julie Siroux,
 ✉ julie.siroux@ext.uca.fr

RECEIVED 06 October 2025
 REVISED 27 April 2026
 ACCEPTED 08 June 2026
 PUBLISHED 25 June 2026

CITATION
 Siroux J, Lemay J, Ramos I, Drapeau V,
 Chaput J-P, Vedrine E, Dalmais C,
 Rousset-Thiery V, Thivel D, Boirie Y and
 Miolanne M (2026) Effects of a family
 home-based intervention on global health
 and 24h movements: an exploratory
 family cluster-based approach to
 childhood obesity.
Int. J. Public Health 71:1609165.
 doi: 10.3389/ijph.2026.1609165

COPYRIGHT
 © 2026 Siroux, Lemay, Ramos, Drapeau,
 Chaput, Vedrine, Dalmais, Rousset-Thiery,
 Thivel, Boirie and Miolanne. This is an
 open-access article distributed under the
 terms of the Creative Commons
 Attribution License (CC BY). The use,
 distribution or reproduction in other
 forums is permitted, provided the original
 author(s) and the copyright owner(s) are
 credited and that the original publication
 in this journal is cited, in accordance with
 accepted academic practice. No use,
 distribution or reproduction is permitted
 which does not comply with these terms.

Effects of a family home-based intervention on global health and 24h movements: an exploratory family cluster-based approach to childhood obesity

Julie Siroux^{1,2*}, Juliette Lemay³, Inès Ramos⁴, Vicky Drapeau⁵,
 Jean-Phillipe Chaput⁶, Elodie Vedrine², Carla Dalmais²,
 Victoria Rousset-Thiery², David Thivel^{1,7}, Yves Boirie^{2,8} and
 Magalie Miolanne^{2,8}

¹Clermont Auvergne University, EA 3533, Laboratory of the Metabolic Adaptations to Exercise Under Physiological and Pathological Conditions (AME2P), CRNH Auvergne, Clermont-Ferrand, France, ²Specialized Center for Obesity in Auvergne (CSO CALORIS), CHU Gabriel Montpied, Clermont-Ferrand, France, ³The Faculty of Agriculture and Food Sciences, School of Nutrition, Laval University, Québec, QC, Canada, ⁴Unit of Biostatistics (DRCI), Clermont-Ferrand University Hospital, Clermont-Ferrand, France, ⁵Department of Kinesiology, Faculty of Educational Sciences, Laval University, Québec, QC, Canada, ⁶Department of Pediatrics, Faculty of Medicine, University of Ottawa, Ottawa, ON, Canada, ⁷International Research Chair Health in Motion, Clermont Auvergne University Foundation, Clermont-Ferrand, France, ⁸Department of Human Nutrition, CHU Gabriel Montpied, Clermont-Ferrand, France

Objectives: This study aimed at (i) exploring the effects of a 12-month home-based family intervention on perceived global health and 24 h movement patterns among children with overweight/obesity (OW/OB) and their family members; (ii) identifying intra-family behavioral clusters and their influence on the intervention's effectiveness.

Methods: 142 families (n = 223 legal guardians) with at least one child with OW/OB were included. At baseline and 12 months, perceived health, sleep, physical activity (PA) and sedentary behaviors (SB) were assessed.

Results: Children with OW/OB showed poorer physical and mental quality of life (QoL) than normal weight (NW) ones (p < 0.001), and decreased body mass index (BMI) z-score over time (p = 0.001). Guardians with OW/OB improved physical QoL at the end of the program (p = 0.002). For all outcomes, clusters analysis suggested a distinction between NW children and those with OW/OB. Children tended to resemble their guardians of same weight status (WS). Changes in BMI z-scores among children with OW/OB might vary by QoL and behavioral profiles with greatest reductions in clusters having higher baseline global health or PA.

Conclusion: This study suggests the potential of family-centered strategies addressing childhood obesity.

KEYWORDS

clusters analysis, home and family-based intervention, movement behaviour, paediatric obesity, perceived health

Introduction

After decades of increasing prevalence, pediatric obesity is now a major threat to future population health. Reports from the World Health Organization (WHO) highlight that over 390 million youth aged 5–19 years were overweight (OW) in 2022, including 160 million living with obesity (OB) [1]. A systematic review and meta-analysis found that 55% of children, and 80% of adolescents with OB will remain affected later in life [2], highlighting the importance of early intervention. Evidence suggests that the factors contributing to childhood obesity often originate at the family level, and that improvements in family lifestyle can positively influence healthy behaviors, helping to reduce the prevalence of childhood obesity [3]. Previous studies suggested an intergenerational transmission of health-related behaviors, with parenting patterns persisting across generations [4–6]. These findings emphasize that childhood obesity is not an isolated issue affecting the child alone but part of a familial context. For example, higher levels of parental physical activity (PA) are associated with increased PA and reduced sedentary behaviors (SB) in children [7], with mothers playing a key role in helping their children meet the 24-h movement guidelines [8]. Similarly, guardians' sleep habits and overall health-related quality of life (HRQoL) are associated with their children's sleep quality and duration [9], as well as with perceived health [10].

Multiple settings and approaches have been proposed for the prevention and management of pediatric obesity, ranging from clinics and schools to surgery and pharmacology. Among these, multidisciplinary behavioral interventions remain crucial for reinforcing and sustaining the benefits achieved [11]. As recently demonstrated, adding behavioral components such as family therapy to diet and PA interventions can positively impact children's body mass index (BMI) and BMI z-scores [11]. Parenting practices promoting healthy behaviors may also be key elements of preventive interventions aimed at reducing the risk of childhood obesity [12]. Given the importance of healthy behaviors in both preventing and managing—and the critical role legal guardians play in shaping their children's lifestyles—family-based interventions involving not only the child but also their guardians and siblings have become increasingly common. These interventions have shown promise in supporting weight management [13] and improving health behaviors [14]. Often implemented successfully in community or care centers [15], family interventions coordinate nutrition, PA, and psychosocial strategies to improve outcomes for children with obesity. However, long-term studies remain rare, and although many interventions adopt family-centered approaches, few incorporate a home-based component that reflects the family's everyday life. Additionally, little is known about the impact of these interventions on family members' health behaviors, and whether distinct intra-family behavioral profiles may influence program success. In this context, the objective is to evaluate the effects of a 12-month family and home-based intervention on perceived health, HRQoL, and

24 h movement patterns (PA, SB, and sleep) among children with OW/OB and their family members. Secondly, the study aims to identify intra-family behavioral clusters and examine their influence on interventions' effectiveness.

Methods

Design

The ProxOb program is a family- and home-based intervention addressing childhood obesity. It engages the family unit and relies on an interdisciplinary team, leveraging parental involvement as a key driver for sustainable lifestyle changes. The program provides tailored advice on PA, SB, nutrition, sleep, psychological and family dynamics. Each family works with a dedicated team of three professionals. Detailed descriptions of the program are available elsewhere [16]. The ProxOb program follows five steps to deliver a personalized intervention: (i) initial assessment: a first meeting with the family, the three professionals (a PA educator, dietician, psychologist, or social worker), the pediatrician, and the program coordinator; (ii) intervention (1 year): a family- and home-based intervention comprising 18 support sessions over 12 months, including three educational assessments, and fifteen workshops (five on nutrition, parenting, and PA); (iii) post intervention assessment: a progress review after the intervention; (iv) maintenance phase (second year): continued support through follow-up calls and online workshops for families; and (iv) long-term assessment: concluding meeting to evaluate the program's impact, with referrals to additional health services if needed for ongoing support. Our study focused on step 1 to 3 with collected data from February 2021 to November 2024 for all family members.

Participants

Families were included through a broad, community-based approach across several French departments (Puy-de-Dôme, Allier, Cantal, Haute-Loire, and Drôme-Ardèche). Inclusion was open, with families entering the program via referrals from healthcare providers or on their own. To participate in the program, families must have at least one child under 18 living with OW/OB, defined as a BMI exceeding the International Obesity Task Force (IOTF) threshold of 25. Additionally, they must meet at least one of the following criteria: (i) be classified as socially vulnerable based on the EPICES score (Évaluation de la Précarité et des Inégalités de santé dans les Centres d'Examens de Santé) [17]; (ii) live in a medically underserved area; (iii) lack access to healthcare services; (iv) have previously failed an obesity intervention. Eligible families were also required to demonstrate sufficient availability to complete the program's questionnaires. The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of the Sud Est VI (2016/CE62). The program was also registered with the Commission National de l'Informatique et des Libertés: 1989437V1. All adult participants completed a written

informed consent form, and a legal guardian signed the form for youth under the age of 18.

Outcomes

All family members (guardians and children, being NW or OW/OB) were evaluated, with the measurement process thoroughly detailed in a previous publication [16]. The following section outlines the specific outcomes used in the study.

Anthropometry

Body height and weight were measured using a standing stadiometer and a body weight scale (Seca Les Mureaux, France) for all participants. In children, BMI, BMI z-scores and percentiles were calculated, with obesity defined using the IOTF age- and sex-specific BMI curves [18]. For adults, BMI was calculated using the formula weight/height^2 (kg/m^2), and weight status (WS) was determined based on WHO reference values [19].

Questionnaires

Adults and children (aged 8–17 years) completed questionnaires covering a range of behaviors and overall health at baseline and after the first year of ProxOb intervention. The study design is presented in Supplementary Material 1.

Health-related quality of life

In adults, Short Form Questionnaire (SF-36), a valid and reliable scale [20] was used to assess HRQoL. It consists of 36 items assessing eight dimensions (physical functioning, physical role, bodily pain, general health, vitality, social functioning, emotions, and mental health), with two final scores (physical and psychological) comprised between 0 (severe dependence) and 100 (independence). For children, HRQoL was assessed using the Pediatric Quality of Life Inventory (PedsQL), a valid questionnaire designed for individuals aged 2–18 years [21]. It consists of 23 questions, divided into four dimensions (physical, emotional, social, and academic) and scored on a 5-point Likert scale ranging from 0 (never) to 4 (almost always) with age-appropriated versions. After recoding, a physical and psychological health scores between 0 and 100 were obtained (scoring method is available in Supplementary Material 2), with higher results indicating higher HRQoL.

Perceived health status

The 3-Levels European Quality of Life 5 Dimensions questionnaire (EQ5D-3L), shown to exhibit excellent psychometric properties in people with NW and with OW/OB was used to assess health status in adults [22]. Is based on five dimensions (mobility, autonomy, usual activities, pain/discomfort, and anxiety/depression), each with three levels (no problems, some problems, and extreme problems). Participants rated their health status by selecting the statement that best described their condition. Scores ranged from 1 to 3, with the mean score calculated across

dimensions. Higher scores indicated poorer health status. In children over 8 years old, health status was assessed using the 3-Levels European Quality of Life 5 Dimension for Youth (EQ5D-3L-Y), an adapted version of the EQ5D-3L. It contains the same five dimensions formulated in an appropriate way for children (mobility, taking care of myself, doing my everyday activities, pain/discomfort, worried/sad/unhappy), with the same scoring method. This questionnaire has been validated and shown to be reliable and feasible for use in young people with health disorders. Its reliability has also been confirmed in a systematic review of 40 studies, in which the mean age of participants was 11.8 years [23].

Sleep

To assess sleep duration, participants were asked “*Usually, how much sleep do you get per night during the week?*” and “*Usually, how much sleep do you get per night at the weekend or on holiday?*.” The average of the two responses was then calculated to assign a score between 0 and 3 to sleep duration, with higher sleep duration equating to higher score. For adults, the classification was: < 5 h = 0, 5–7 h = 1, 7–9 h = 2 and > 9 h = 3. For children, the categories were as follows: <7 h = 0, 7–9 h = 1, 9–11 h = 2, and > 11 h = 3. This classification was made according to the sleep duration recommendations for both pediatric [24] and adult populations [25].

Physical activity and sedentary behaviors

To obtain indicators of PA and SB, adults were asked to complete the Physical Activity Questionnaire of the National Observatory on Physical Activity and Sedentarity (ONAPS-PAQ), a 21-question questionnaire in three parts, covering work, transport, and leisure/home activities during a usual week [26]. This reliable and valid questionnaire in adults aged 18–69 years [26] gives a score for the level of PA and the amount of sedentary time, with higher scores indicating better PA level and higher sedentarity. Children aged 8 years or older filled in the Children and Adolescents Physical Activity and Sedentarity Questionnaire (CAPAS-Q), which enables a similar assessment [27]. This questionnaire has also been validated in children and adolescents aged 8–18 years [27]. Using different contexts, it also provides two scores (PA and SB). The scoring method and interpretation are available in Supplementary Material 2.

Statistical analyses

Statistical analyses were performed using Stata software (version 15; StataCorp, College Station, TX, USA). All tests were two-sided, with an alpha level set at 5%. Continuous data were presented with mean \pm standard deviation and categorical data with number and associated proportions. Longitudinal analyses were performed using mixed linear regression models to assess the effect of the intervention (T0 vs. T12), WS (NW vs. OW/OB) and sex (female vs. male) as fixed covariates, and considering individuals and family as random effects, on health status, sleep, PA and SB. These models were conducted on legal guardians and children separately.

TABLE 1 Evolution of body mass index z-score, health-related quality of life, health status and movement behaviors (sleep, physical activity, sedentary behaviors) in children (Auvergne, France, 2025).

	All children			NW children		Children with OW/OB		Mixed Models effects NW vs. OW/OB		
	T0 n = 254	T12 n = 73	Time	T0 n = 60	T12 n = 16	T0 n = 194	T12 n = 57	Time	Weight status	Time x WS
BMI z-score IOTF ¹	2.1 ± 1.1	2.1 ± 1.0	0.074	0.6 ± 0.7	0.9 ± 0.3	2.5 ± 0.8	2.3 ± 0.9	0.044	<0.001	0.001^a
Physical health (HRQoL)	79.8 ± 13.5	80.2 ± 14.6	0.548	86.8 ± 13.1	83.9 ± 4.2	78.0 ± 13.1	79.4 ± 16.0	0.615	<0.001	0.422
Psychological health (HRQoL)	73.8 ± 14.5	73.5 ± 11.7	0.815	82.3 ± 12.4	74.7 ± 8.0	71.8 ± 14.3	73.2 ± 12.4	0.293	<0.001	0.191
Health status	1.3 ± 0.3	1.3 ± 0.3	0.300	1.1 ± 0.2	1.2 ± 0.2	1.3 ± 0.3	1.3 ± 0.3	0.680	<0.001	0.350
Sleep duration (hrs)	9.5 ± 1.0	9.5 ± 1.0	0.167	9.3 ± 1.0	9.7 ± 0.9	9.5 ± 1.0	9.4 ± 1.0	0.852	0.460	0.387
Sleep duration^{2, 3}										
0	2 (2)	0 (0)		1 (4)	0 (0)	1 (1)	0 (0)			
1	28 (25)	11 (28)		7 (29)	1 (12)	21 (24)	10 (32)			
2	77 (68)	25 (64)		15 (63)	6 (76)	62 (70)	19 (61)			
3	6 (5)	3 (8)		1 (4)	1 (12)	5 (5)	2 (7)			
Physical activity level	2.2 ± 0.4	2.2 ± 0.4	0.739	2.2 ± 0.3	2.2 ± 0.5	2.2 ± 0.4	2.1 ± 0.3	0.226	0.942	0.249
Sedentary behaviors level	2.6 ± 0.7	2.7 ± 0.8	0.086	2.6 ± 0.7	2.6 ± 0.9	2.6 ± 0.7	2.7 ± 0.8	0.633	0.930	0.704

Abbreviations. BMI: body mass index, IOTF: international obesity task force, HRQoL: Health-Related Quality of Life, hrs: hours, OB: obesity, OW: overweight, T0: baseline, T12: after 1-year intervention, WS: weight status.

Sex effect was observed for sedentary behaviors, Sex effect was observed for sedentary behaviors (p = 0.008), data not shown for children, contrary to legal guardians for which sex differences are well known and of interest. P-values in bold are < 0.05.

¹Quantitative variables: Mean ± SD, with significant p-value < 0.05.

²Sleep duration divided into 4 categories, as described in Methods.

³Qualitative variables: n (% in the subsample).

^aSignificant difference between children with OW/OB, at T0 and children with OW/OB, at T12 (p = 0.006).

Interaction between time and WS was assessed in a complementary model. A Sidak's type I error correction was applied for multiple pairwise comparisons.

A sensitivity (complete-case) and attrition analysis comparing baseline characteristics between participants with and without T12 data were performed. Model assumptions were checked using residual diagnostics, including assessment of normality and homoscedasticity, and no major violations were observed. Mixed models were estimated using maximum likelihood, allowing inclusion of participants with incomplete data under a missing-at-random assumption. Spearman's correlations were used to measure the association between BMI z-score change and changes in health status, sleep, PA and SB (Δvalue = value T12 - value T0). Correlation coefficients were interpreted as follows (absolute coefficient value): ≥ 0.70 (strong), 0.40 to 0.69 (moderate), 0.10 to 0.39 (weak), < 0.10 (negligible). To classify participants by their initial characteristics and behavior at baseline, a principal component analysis (PCA) was performed in children, using FactoMineR and Factoextra packages. A multiple component analysis was also performed, categorizing the scores using

quartile 1, median and quartile 3 as cut-offs, for children and legal guardians separately. These categories allowed to consider children and adults on the same classification. Hierarchical clustering on principal components was then performed. Cluster analyses were performed using complete case and are considered exploratory. Finally, BMI z-scores were compared between T0 and T12 in each cluster using a paired Wilcoxon test.

Results

One hundred forty-two families were included in the analysis, representing 477 participants (135 female guardians, 88 male guardians, 60 NW children, and 194 children with OW/OB). Regarding the WS of legal guardians, 79% (n = 107) of females and 85% (n = 75) of males were living with OW/OB (Supplementary Material 3). Sex parity was almost observed in children with OW/OB (with 53% of females and 47% of males) and in NW children (50% of both females and males) All the descriptive characteristics are shown in Supplementary Material 3. The attrition analysis comparing

TABLE 2 Evolution of body mass index, health-related quality of life, health status and movement behaviors (sleep, physical activity, sedentary behaviors) in legal guardians (Auvergne, France, 2025).

	Female legal guardians						Male legal guardians				Mixed models effects females vs. males				
	All legal guardians			NW		OW/OB		NW		OW/OB		Sex	Time	WS	Time x WS
	T0 n = 223	T12 n = 99	Time	T0 n = 28	T12 n = 11	T0 n = 107	T12 n = 57	T0 n = 13	T12 n = 4	T0 n = 75	T12 n = 27				
BMI (kg/m ²) ¹	32.8 ± 8.1	32.2 ± 5.5	0.902	22.6 ± 1.6	24.5 ± 3.7	35.9 ± 8.1	32.9 ± 5.1	23.7 ± 1.2	25.2 ± 0.6	33.8 ± 5.9	34.1 ± 4.4	0.071	0.033	<0.001	0.017^a
Physical health (HRQoL)	71.3 ± 17.8	78.3 ± 17.8	<0.001	84.8 ± 8.4	80.3 ± 12.6	66.7 ± 17.1	75.3 ± 16.3	86.3 ± 3.8	86.7 ± 5.5	72.1 ± 19.3	82.0 ± 22.5	0.079	0.252	<0.001	0.002^b
Psychological health (HRQoL)	60.1 ± 18.5	63.4 ± 19.9	0.081	64.0 ± 13.1	61.5 ± 18.7	56.4 ± 18.7	61.4 ± 19.6	70.9 ± 15.0	66.1 ± 10.0	62.4 ± 19.3	68.2 ± 22.0	0.008	0.506	0.035	0.110
Health status	1.4 ± 0.3	1.4 ± 0.3	0.021	1.2 ± 0.1	1.3 ± 0.2	1.4 ± 0.3	1.4 ± 0.3	1.2 ± 0.1	1.2 ± 0.2	1.4 ± 0.3	1.3 ± 0.3	0.126	0.683	<0.001	0.571
Sleep duration (hrs)	8.0 ± 0.8	8.1 ± 1.0	0.115	8.1 ± 0.9	8.1 ± 1.0	8.1 ± 0.9	8.3 ± 1.1	7.8 ± 0.7	7.3 ± 0.4	7.8 ± 0.8	7.9 ± 0.8	0.002	0.894	0.627	0.438
Sleep duration ^{2, 3}															
Very low	0 (0)	0 (0)		0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)				
Low	13 (10)	7 (10)		2 (13)	1 (12)	4 (6)	4 (9)	2 (22)	0 (0)	5 (12)	2 (11)				
Moderate	94 (72)	45 (64)		8 (53)	5 (63)	48 (74)	25 (60)	6 (67)	2 (100)	31 (76)	13 (72)				
High	24 (18)	18 (26)		5 (34)	2 (25)	13 (20)	13 (31)	1 (11)	0 (0)	5 (12)	3 (17)				
Physical activity (METs.min ⁻¹ .week ⁻¹)	96.0 ± 150.9	88.1 ± 115.7	0.753	80.9 ± 92.3	50.1 ± 59.2	77.0 ± 135.6	65.2 ± 91.3	294.9 ± 427.7	221.4 ± 282.0	101.1 ± 108.4	134.9 ± 140.7	0.016	0.570	0.109	0.404
Sedentary time (hrs)	7.6 ± 6.3	7.6 ± 4.8	0.755	7.3 ± 5.5	10.1 ± 7.8	8.0 ± 7.5	7.3 ± 3.6	5.8 ± 4.9	4.3 ± 2.3	7.5 ± 4.8	7.7 ± 5.4	0.538	0.194	0.386	0.191


Abbreviations. BMI: body mass index, HRQoL: Health-Related Quality of life, hrs: hours, NW: normal weight, OB: obesity, OW: overweight, T0: baseline, T12: after 1-year intervention, WS: weight status.

¹Quantitative variables: Mean ± SD, with significant p-value <0.05. ²Sleep duration divided into 4 categories, as the described in Methods. P-values in bold are < 0.05.

³Qualitative variables: n (% in the subsample).

^aNo significant differences in *post hoc* analysis.

^bSignificant difference between guardians with OW/OB, at T0 and guardians with OW/OB, at T12 (p < 0.001).



baseline characteristics between completers and non-completers only showed a significant age difference for children ($p = 0.022$), with no difference observed for adults (Supplementary Materials 4,5).

Effects of the ProxOb program at the individual level depending on weight status

Tables 1, 2 display all the detailed results for children (Table 1) and adults (Table 2) regarding the effects of the ProxOb program on HRQoL, PA, SB, and sleep at the individual level. Regarding children, results for BMI z-scores, physical and psychological HRQoL, and perceived health status were significantly different between children with OW/OB and their NW counterparts who shown lower BMI z-scores, higher HRQoL and perceived health status (all p -values <0.001) (Table 1), as also found by clusters analyses (Supplementary Materials 6, 7). A significant time \times WS interaction was also found for BMI z-score, with the *post hoc* analysis showing significant decrease from T0 (2.5 ± 0.8) to T12 (2.3 ± 0.9) among children with OW/OB ($p = 0.006$). A significant time effect in BMI z-score was observed in children ($p = 0.044$), with a corresponding increase in NW children ($\beta = 0.236$, 95% CI [0.010, 0.470]) (effect sizes are presented in Supplementary Material 8). Finally, as shown in Supplementary Material 9, no significant correlation was observed between the delta changes in each behavior and the delta BMI z-score of children with OW/OB,

between baseline and 1 year of intervention. Specifically, correlation deltas ranged from 0.146 (Δ health status) to -0.500 (Δ PA) and p -values from 0.132 (Δ physical health) to 0.957 (Δ SB).

In adults, results showed significantly higher BMI ($p < 0.001$), and lower health status ($p < 0.001$), physical ($p < 0.001$) and psychological ($p = 0.035$) HRQoL in individuals with OW/OB compared to NW. Sex differences were observed for psychological health ($p = 0.008$), sleep duration ($p = 0.002$), and PA ($p = 0.016$) with females reporting longer sleep and males having a higher perceived psychological health and PA level. Regarding physical HRQoL, a time \times WS interaction was observed ($p = 0.002$), with improved perceived physical health in guardians with OW/OB at T12 (females: 75.3 ± 16.3 ; males: 82.0 ± 22.5) compared to baseline (66.7 ± 17.1 ; 72.1 ± 19.3 ; $p < 0.001$) (Table 2). As in children, a significant time effect on BMI was observed in guardians ($p = 0.033$), with a corresponding increase ($\beta = 1.400$, 95% CI [0.110, 2.700]) (all effect sizes presented in Supplementary Material 10).

Importantly, sensitivity analyses yielded similar results for BMI and physical health in guardians, while for children, results were only similar for BMI z-score IOTF, which showed significant time, WS, and interaction effects. Details are provided in Supplementary Material 11 (children) and Supplementary Material 12 (guardians), and data availability in Supplementary Material 13.

Baseline associations between family members

The graphical results of multi-component analysis provide a visual representation of the patterns and associations with families for HRQoL (Figure 1). When considering all family members, the three clusters of HRQoL were widely different, as shown in Table 3, with cluster 3 indicating a better health. This cluster included almost all of the NW children (80%), while those with OW/OB were mostly in clusters 1 and 2, indicating poorer HRQoL. In general, children with OW/OB were close to their legal guardians of same WS, with cluster 1 including almost only guardians and children with OW/OB. Cluster 2 tended to be more heterogeneous, as it also comprised most individuals with OW/OB and almost all NW female legal guardians (69%), suggesting that most females had higher HRQoL than their male counterparts, despite low sample sizes.

Although clustering was similar for sleep, PA, and SB, NW female legal guardians and children were all grouped in cluster 3, whereas adults and children with OW/OB were mainly in cluster 1 (Table 4).

As shown in Figure 2, children generally clustered with legal guardians of same WS (Figure 2). Finally, in contrast to clusters 1 and 3, cluster 2 included more males, with higher PA level, and lower SB.

Baseline behaviors profile and success of the intervention

Changes in BMI z-scores among children with OW/OB between baseline and end of intervention are presented by cluster in Supplementary Materials 6, 7 (children) and Tables 3, 4 (all

TABLE 3 Identification of three distinct family clusters based on health-related quality of life and health status using a multi-component analysis: distribution of family members by sex and weight status (Auvergne, France, 2025).

	Cluster 1 – Low global health (vulnerable profile) n = 76	Cluster 2 – Moderate global health (intermediate profile) n = 155	Cluster 3 – High global health (favourable profile) n = 62
Physical health (HRQoL) ¹			
Low physical health	68 (90)	20 (13)	0 (0)
Moderate physical health	7 (9)	117 (75)	11 (18)
High physical health	1 (1)	18 (12)	51 (82)
Psychological health (HRQoL)			
Low psychological health	54 (71)	28 (18)	0 (0)
Moderate psychological health	20 (26)	110 (71)	15 (24)
High psychological health	2 (3)	17 (11)	47 (76)
Health status			
High health status	2 (3)	13 (8)	49 (79)
Moderate health status	18 (24)	134 (87)	13 (21)
Low health status	56 (74)	8 (5)	0 (0)
Sex and weight status			
Female legal guardians NW	0 (0)	11 (7)	5 (8)
Female legal guardians OW/OB	23 (30)	49 (32)	4 (7)
Male legal guardians NW	0 (0)	4 (2)	3 (5)
Male legal guardians OW/OB	20 (26)	17 (11)	5 (8)
Children NW	3 (4)	3 (2)	23 (37)
Children OW/OB	30 (40)	71 (46)	22 (35)
Δ BMI z-score, children with OW/OB ²	-0.2 ± 0.4	-0.1 ± 0.2*	-0.5 ± 0.7

Abbreviations. BMI: body mass index, HRQoL: Health-Related Quality of life, NW: normal weight, OB: obesity, OW: overweight, Δ BMI z-score: z-score change from baseline (T0) to the end of the intervention (T12).

¹n (% in the cluster).

²Mean ± SD.

*p-value between T0 and T12 < 0.05.

³category classification based on a quartile analysis of the scores as follows: low: ≤ Q1, moderate: > Q1 and < Q3, high: ≥ Q3. The cut-offs used for adults are: physical health Q1 (low) = 63.2/Q3 (high) = 87.7, psychological health Q1 (low) = 47.1/Q3 (high) = 75.7, and health status Q1 (high) = 1.2/Q3 (low) = 1.6. For children the cut-off values are: physical health Q1 (low) = 68.8/Q3 (high) = 90.6, psychological health Q1 (low) = 65.0/Q3 (high) = 83.3, health status Q1 (high) = 1.0/Q3 (low) = 1.4.

family members). In children clusters, a significant decrease in BMI z-scores was observed in cluster 2 for HRQoL (-0.3 ± 0.4 [-1.4, 0.3], p = 0.003) (Supplementary Material 6) and in cluster 1 for 24-h movement (-0.3 ± 0.4 [-1.4, 0.5], p = 0.026) (Supplementary Material 7). In family clusters, a significant decrease in BMI z-scores was observed in cluster 2 for HRQoL and health status (-0.1 ± 0.2, p = 0.014) (Table 3) while no change was found for sleep, PA, and SB (Table 4).

Discussion

Although various interventions exist for childhood obesity prevention and management, incorporating behavioral components and involving the whole family appear key to their effectiveness and sustainability [4, 11]. Leveraging the ProxOb program, this study examined the effects of a home and family-based intervention on HRQoL and movement behaviors (PA, SB,

TABLE 4 Identification of three distinct family clusters based on sleep duration, physical activity and sedentary behaviors using a multi-component analysis: distribution of family members by sex and weight status (Auvergne, France, 2025).

	Cluster 1 – Sedentary & inactive profile (unhealthy behaviors) n = 105	Cluster 2 – Active profile (healthy behaviors) n = 37	Cluster 3 – Intermediate risk profile n = 26
Sleep duration ¹			
Low sleep duration	29 (28)	6 (16)	9 (35)
Moderate sleep duration	43 (41)	23 (62)	13 (50)
High sleep duration	33 (31)	8 (22)	4 (15)
Physical activity			
Low physical activity level	35 (33)	0 (0)	5 (19)
Moderate physical activity level	69 (66)	6 (16)	13 (50)
High physical activity level	1 (1)	31 (84)	8 (31)
Sedentary behaviors			
Low sedentary behaviors	28 (27)	10 (27)	6 (23)
Moderate sedentary behaviors	46 (44)	21 (57)	13 (50)
High sedentary behaviors	31 (29)	6 (16)	7 (27)
Sex and weight status			
Female legal guardians NW	0 (0)	0 (0)	11 (42)
Female legal guardians OW/OB	40 (38)	5 (13)	0 (0)
Male legal guardians NW	0 (0)	5 (13)	0 (0)
Male legal guardians OW/OB	18 (17)	15 (41)	0 (0)
Children NW	0 (0)	0 (0)	15 (58)
Children OW/OB	47 (45)	12 (32)	0 (0)
Δ BMI z-score children OW/OB ²	-0.1 ± 0.4	-0.4 ± 0.7	-

Abbreviations. BMI: body mass index, NW: normal weight, OB: obesity, OW: overweight, Δ BMI z-score: z-score change from baseline (T0) to the end of the intervention (T12).

¹n (% in the cluster).


²Mean ± SD.

³category classification based on a quartile analysis of the scores as follows: low: ≤ Q1, moderate: > Q1 and < Q3, high: ≥ Q3. The cut-offs used for adults are: sleep duration Q1 (low) = 7.5/Q3 (high) = 8.8, physical activity Q1 (low) = 9.3/Q3 (high) = 129.0, and sedentary behaviors Q1 (low) = 3.3/Q3 (high) = 9.8. For children the cut-off values are: sleep duration Q1 (low) = 8.9/Q3 (high) = 10.3, physical activity Q1 (low) = 1.9/Q3 (high) = 2.4, sedentary behaviors Q1 (low) = 2.2/Q3 (high) = 3.2.

and sleep). These effects were analyzed at the individual family-member level, and an exploratory cluster analysis was conducted to explore intra-family associations. This study also investigated whether baseline associations among family members might be related to the intervention' success. The findings support the potential effectiveness of the ProxOb program in reducing BMI z-scores in children with OW/OB, although no significant time effects were observed for individual behaviors in mixed-model. Clusters exploratory analyses also revealed distinct patterns of HRQoL and movement behaviors, with similarities in HRQoL and perceived health status across weight categories. Specifically, better physical and psychological health perceptions was associated with improved health status in NW participants. In terms of

movement behaviors, higher levels of PA seem to be associated with lower SB, while sleep did not emerge as a key factor in clustering.

Firstly, the observed changes in BMI z-scores align with a previous systematic review and meta-analysis showing that 73% of home-based interventions positively impacted BMI z-scores [28], with reductions up to 37% across 22 home-based interventions [28]. The individual level mixed-model analysis also indicates lower HRQoL in children with OW/OB compared with NW peers, a result consistent with the findings of Van de Pas and colleagues [29]. Consistent with such findings, family-based obesity prevention programs may be among the most effective strategies for targeting childhood obesity and reducing BMI z-scores [30].



Despite differences in WS, our results do not show any effect of the intervention on children's HRQoL or health status. This lack of intervention effect may partly be explained by the high scores at baseline, a caveat to consider when interpreting the results, as already outlined in literature [31]. Among adults, an effect was observed for physical HRQoL, that significantly improved in guardians with OW/OB post-intervention. This result is in line with a previously conducted 6-month family intervention, with an improvement of the guardians' general health perceptions [32], although the questionnaire administered was different from the SF-36 used in the present study. Interestingly, this observed improvement occurred in a cohort of adults with a mean baseline BMI of 35.6 kg/m², highlighting the potential cumulative effects of family obesity interventions on guardians with OW/OB [32].

Regarding movement behaviors, none was modified in response to the ProxOb intervention in children and adults. Interestingly, these findings align with the conclusions of a systematic and meta-analytic review, highlighting that family-based childhood obesity interventions have limited effects on children's and guardians' movement behaviors [33]. These results may be explained by practical barriers such as time constraints, competing priorities or unrealistic expectations [34]. Resources limitation and limited access to supportive environments may also have contributed, particularly given the program's inclusion of socially vulnerable and rural family [35]. Finally, in line with recent studies, the interplay of family dynamics, including cohesion [36],

intergenerational influences [37] and adaptive capacity [37], may play a key role in the success of family-based interventions, although these factors were not assessed in the present study.

Interestingly, exploratory cluster analyses showed concordance between legal guardians and children of same WS regarding HRQoL and movement behaviors, in line with prior evidence linking guardian to child health [10] and PA [38]. The intergenerational transmission of obesity and obesity-related behaviors are mediated by genetic and environmental factors, with guardians playing a key role in shaping children's behaviors [39]. Importantly, this clustering between family members appeared to be higher between children and their mothers, consistent with previous literature highlighting that mothers may have a significant influence on their children's health and wellbeing [40], but also on PA and SB [41, 42].

Furthermore, as shown by the WS effects analyses of the linear mixed models, our exploratory cluster analysis among children suggested differences by WS, with most NW children being in the same cluster and reporting better perceived HRQoL and health status than their counterparts with OW/OB, in line with previous research showing lower HRQoL scores in children with OW/OB compared to their NW peers [43, 44]. Children with OW/OB also showed reduced psychological health, self-esteem, and physical functioning [44], with such effects observed in both clinical and community-based samples [43], highlighting the importance of considering HRQoL dimensions in obesity prevention and management interventions, as excess weight affects not only biological health, but also overall HRQoL.

With regard to movement behaviors, children were found divided according to their WS, but the results were more heterogeneous, with no difference between children with NW and those with OW/OB. However, studies using subjective questionnaires suggest that children, regardless of WS, report similar sleep duration, PA levels, and SB [45]. This highlights the limitations of self-reported measures, which may partly explain the present findings. In all family member clusters, the observation that higher PA levels may be linked with lower sedentarity highlights the importance of the interplay between PA practice and the reduction of SB.

Finally, although no significant correlations were found between individual health behaviors and changes in BMI z-scores, the exploratory cluster analysis indicated greater reductions in BMI z-scores among children in clusters characterized by better perceived health and higher PA levels. These findings suggest that positive health perceptions and regular PA may serve as key levers for intervention success, potentially enhancing participant engagement and overall effectiveness. This aligns with previous studies showing that children who were already physically active before an intervention experienced the greatest benefits in terms of weight reduction [46]. These results also highlight the complexity of obesity and the fact that others behavioral changes (e.g., eating habits) may be associated with BMI z-scores changes and intervention success. To our knowledge, no prior study has reported a similar association between baseline HRQoL and reductions in BMI z-scores.

All of the present findings should be interpreted in light of certain limitations. These include the subjective nature of measuring PA, SB, and sleep duration—despite the use of validated

questionnaires. Additionally, the high rate of missing data limited the scope of the analyses, making it impossible to conduct principal component or multi-component analyses and thereby restricting the ability to interpret the results in cluster form. This also limits the robustness of clustering-based interpretation given the sample size and missing data constraints. The relatively high attrition rate and thus reduced sample size at follow-up, particularly in subgroup and cluster analyses, may also limit statistical power and robustness of these findings. As such, cluster analyses should be considered exploratory and interpreted with caution. Importantly, this attrition rate includes both families who left the program for personal and/or organizational reasons and families for whom data are missing (who did not leave the clinical program) but did not complete the assessments. This should prompt us to reflect on this type of intervention, considering both the organizational aspects and the type of assessment implemented, which should not be too burdensome for families. It would also have been valuable to assess participants' readiness to change, as this has been shown in other studies to influence intervention outcomes and could have provided further insights. Similarly, family functioning and the context of psychosocial complexity were not analyzed in the present study. With larger sample sizes, it might be interesting to carry out analyses according to the age group of the children and the extent of their excess weight. Importantly, it should be noted that while the ProxOb program offers valuable, innovative, and real-world insights, it was not originally designed as a scientific study. Instead, it is a field-based intervention aimed at preventing and treating childhood obesity at the family level. Although this limits the strength of the conclusions that can be drawn, it also highlights the practical feasibility of implementing large-scale, family-based, home-centered programs—providing unique evidence grounded in real-world practice.

Conclusion

By integrating behavioral and multidisciplinary approaches, the ProxOb program suggests associations between the participation in the intervention and reduced BMI z-scores in children with OW/OB, as well as increased physical HRQoL in adults with OW/OB. The present study also indicates the potential existence of distinct behavioral phenotypes related to HRQoL, PA, SB, and sleep within family members. These findings reinforce the importance of family-based interventions in managing childhood obesity and support the relevance of considering intergenerational transmission of health behaviors. This study particularly suggests that children's health behaviors may be aligned with those of their legal guardians, with family members displaying similar behavioral profiles based on sex and WS, underscoring the important influence of legal guardians in shaping children's lifestyle habits. The results also indicate that positive health perceptions and high levels of PA could be key factors in enhancing intervention success among children, supporting the need for personalized strategies that consider individual and family behavioral patterns. Overall, this study provides valuable insights into associations between a family and home-based intervention and health-related outcomes in the context of childhood obesity, while also emphasizing the added value of conducting family behavioral profiling in future interventions. Further research should explore the scalability of such approaches, including a

better understanding of implementation aspects and the overall costs associated with delivering multi-professional, long-term family-based programs, in order to inform their potential replication in broader settings.

Data availability statement

Data supporting these findings are available from the corresponding author upon request. The data are not publicly available due to privacy or ethical restrictions.

Ethics statement

The studies involving humans were approved by Ethics Committee of the Sud Est VI Ethics Committee (2016/CE62). The studies were conducted in accordance with the local legislation and institutional requirements. Written informed consent for participation in this study was provided by the participants' legal guardians/next of kin.

Author contributions

Conceptualization: MM and YB; Methodology: MM and YB; Investigation: EV and CD; Formal analysis: IR, DT, and JS; Data management: EV, CD, JS, and VR; Writing JS, JL, DT, and MM; Review and editing: MM, VD, J-PC, and YB. All authors contributed to the article and approved the submitted version.

Funding

The author(s) declared that financial support was received for this work and/or its publication. This work was supported by the Regional Sanitary Agency Auvergne/AuRA and National Health Insurance Fund.

Acknowledgements

The authors would like to express their gratitude to all the practitioners who were involved in creating, developing and improving the ProxOb project. They would also like to thank all the families who participated in the program.

Conflict of interest

The authors declare that they do not have any conflicts of interest.

Generative AI statement

The author(s) declared that generative AI was not used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

References

- Phelps NH, Singleton RK, Zhou B, Heap RA, Mishra A, Bennett JE, et al. Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222 million children, adolescents, and adults. *Lancet* (2024) 403(10431):1027–50. doi:10.1016/S0140-6736(23)02750-2
- Simmonds M, L O CG, N. W. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* (2016) 17(2):95–107. doi:10.1111/obr.12334
- Gray LA, Hernandez Alava M, Kelly MP, Campbell MJ. Family lifestyle dynamics and childhood obesity: evidence from the millennium cohort study. *BMC Public Health* (2018) 18(1):500. doi:10.1186/s12889-018-5398-5
- Chen G, Olsen JA, Lamu AN. The influence of parents' and partner's education on own health behaviours. *Soc Sci Med* (2024) 343:343. doi:10.1016/j.socscimed.2024.116581
- Classen TJ. Measures of the intergenerational transmission of body mass index between mothers and their children in the United States, 1981–2004. *Econ Hum Biol* (2010) 8(1):30–43. doi:10.1016/j.ehb.2009.11.002
- Goode A., Mavromaras K, Smith M. *Intergenerational Transmission of Healthy Eating Behaviour and the Role of Household Income*. Bonn, Germany: Institute for the Study of Labor (2008). Available online at: <https://hdl.handle.net/10419/34802> (Accessed August 25, 2025).
- Fuemmeler BF, Anderson CB, Mäse LC. Parent-child relationship of directly measured physical activity. *Int J Behav Nutr Phys Activity* (2011) 8:17. doi:10.1186/1479-5868-8-17
- Sigmundová D, Vorlíček M, Voráčková J, Dyrgrín J, Sigmund E. Parental impact on adherence of young children to 24-h movement behaviour guidelines: the Czech FAMILY physical activity, sedentary behaviour and sleep study. *Eur J Public Health* (2025) 35(2):295–301. doi:10.1093/eurpub/ckae224
- Chehri A, Taheri P, Khazaei H, Jalali A, Ahmadi A, Mohammadi R. The relationship between parents' sleep quality and sleep hygiene and preschool children' sleep habits. *Sleep Sci* (2022) 15(3):272–8. doi:10.5935/1984-0063.20220051
- Musetti A, Zagaria A, Pezzi M, Fante C, Dioni B, Raffin C, et al. Parental quality of life, child adjustment and adult attachment in parents of children and adolescents with autism spectrum disorder. *Res Dev Disabil* (2024) 146:104684. doi:10.1016/j.ridd.2024.104684
- Salam RA, Padhani ZA, Das JK, Shaikh AY, Hoodbhoy Z, Jeelani SM, et al. Effects of lifestyle modification interventions to prevent and manage child and adolescent obesity: a systematic review and meta-analysis. *Nutrients* (2020) 12:1–23. doi:10.3390/nu12082208
- Campos AP, Robles J, Matthes KE, Alexander RC, Goode RW. Parenting practices to prevent childhood obesity among Hispanic families: a systematic literature review. *Child Obes* (2024) 21:113–28. doi:10.1089/chi.2024.0311
- Fernández-Lázaro D, Celorrio SMAM, Garrosa E, Fernández-Araque AM, Mielgo-Ayuso J, Roche E, et al. Evaluation of family-based interventions as a therapeutic tool in the modulation of childhood obesity: a systematic review. *Child Multidisciplinary Digital Publishing Inst (MDPI)* (2024) 11:930. doi:10.3390/children11080930
- Tamayo MC, Dobbs PD, Pincú Y. Family-centered interventions for treatment and prevention of childhood obesity in Hispanic families: a systematic review. *J Community Health* (2021) 25:635–43. doi:10.1007/s10900-020-00897-7
- Anderson YC, Wynter LE, Grant CC, Cave TL, Derraik JGB, Cutfield WS, et al. A novel home-based intervention for child and adolescent obesity: the results of the whānau pakari randomized controlled trial. *Obesity* (2017) 25(11):1965–73. doi:10.1002/oby.21967
- Miolanne M, Lambert C, Masurier J, Cardenoux C, Fillion A, Beraud S, et al. Designing, implementing, and evaluating a home-based, multidisciplinary, family-centered pediatric obesity intervention: the ProxOb program. *Children* (2022) 9(5):737. doi:10.3390/children9050737
- Guéguen R SC the E working group: ALRGFGCLREMPMERBRM. *The EPICES Score: An Individual Index of Material and Social Deprivation Related to Health Status* (2005).
- Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* (2012) 7(4):284–94. doi:10.1111/j.2047-6310.2012.00064.x
- World Health Organization (WHO). *Obesity: Preventing and Managing the Global Epidemic: Report of a WHO Consultation*. Geneva, Switzerland: World Health Organization (2000).

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.sspH-journal.org/articles/10.3389/ijph.2026.1609165/full#supplementary-material>

- Leplège A, Ecosse E, Verdier A, Perneger TV. The French SF-36 health survey: Translation, cultural adaptation and preliminary psychometric evaluation. *J Clin Epidemiol* (1998). 51:1013–1023. doi:10.1016/s0895-4356(98)00093-6
- Tessier S, Vuillemin A, Lemelle JL, Briançon S. Propriétés psychométriques du questionnaire générique français « Pediatric Quality of Life Inventory Version 4.0 » (PedsQLTM 4.0). *Revue européenne de Psychol appliquée* (2009) 59(4):291–300. doi:10.1016/j.erap.2009.06.001
- Feng YS, Kohlmann T, Janssen MF, Buchholz I. Psychometric properties of the EQ-5D-5L: a systematic review of the literature. *Qual Life Res* (2021) 30(3):647–73. doi:10.1007/s11136-020-02688-y
- Golicki D, Młyńczak K. Measurement properties of the EQ-5D-Y: a systematic review. *Value in Health* (2022) 25(11):1910–21. doi:10.1016/j.jval.2022.05.013
- Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, et al. Recommended amount of sleep for pediatric populations: a consensus statement of the American academy of sleep medicine. *J Clin Sleep Med* (2016) 12(6):785–6. doi:10.5664/JCSM.5866
- Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, Buysse D, et al. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American academy of sleep medicine and sleep research society. *Sleep* (2015) 38(6):843–4. doi:10.5665/SLEEP.4716
- Charles M, Thivel D, Verney J, Isacco L, Husu P, Vähä-Yppä H, et al. Reliability and validity of the ONAPS physical activity questionnaire in assessing physical activity and sedentary behavior in French adults. *Int J Environ Res Public Health* (2021) 18(11):5643. doi:10.3390/ijerph18115643
- Fillon A, Pereira B, Vanhelst J, Baran J, Masurier J, Guirado T, et al. Development of the children and adolescents physical activity and sedentary questionnaire (CAPAS-Q): psychometric validity and clinical interpretation. *Int J Environ Res Public Health* (2022) 19(21):13782. doi:10.3390/ijerph192113782
- Pamungkas RA, Chamroonsawasdi K. Home-based interventions to treat and prevent childhood obesity: a systematic review and meta-analysis. *MDPI Multidisciplinary Digital Publishing Inst* (2019) 9:38. doi:10.3390/bs9040038
- Van De Pas KGH, De Krom MAP, Winkens B, Van Dielen FMH, Vreugdenhil ACE. Health-related quality of life in children and adolescents with overweight, obesity, and severe obesity: a cross-sectional study. *Obes Facts* (2023) 16(3):282–92. doi:10.1159/000529560
- Gruber KJ, Haldeman LA, Cdc for. Using the family to combat childhood and adult obesity (2009). Available online at: http://www.cdc.gov/pcd/issues/2009/jul/08_0191.htm (Accessed August 25, 2025).
- Kleppang AL, Abildsnes E, Haraldstad K, Stea TH. Changes in health-related quality of life and sleep habits after a 6-month non-randomised cluster-controlled trial among children with overweight or obesity. *Eur Child Adolesc Psychiatry* (2024) 33(9):3179–87. doi:10.1007/s00787-024-02375-0
- Kalarachian MA, Levine MD, Arslanian SA, Ewing LJ, Houck PR, Cheng Y, et al. Family-based treatment of severe pediatric obesity: randomized, controlled trial. *Pediatrics* (2009) 124(4):1060–8. doi:10.1542/peds.2008-3727
- Arnason A, Langarica N, Dugas LR, Mora N, Luke A, Markossian T. Family-based lifestyle interventions: what makes them successful? A systematic literature review. *Prev Med Rep* (2021) 21:101299. doi:10.1016/j.pmedr.2020.101299
- Howie EK, McManus A, Smith KL, Fenner AA, Straker LM. Practical lessons learned from adolescent and parent experiences immediately and 12 months following a family-based healthy lifestyle intervention. *Child Obes* (2016) 12(5):401–9. doi:10.1089/chi.2016.0091
- Andino J, Park-Mroch J, Francis SL, O'Shea AMJ, Engebretsen B, Rice S, et al. A randomized controlled trial of a community-based obesity intervention utilizing motivational interviewing and community resource mobilization for low-income families: study protocol and baseline characteristics. *Contemp Clin Trials* (2022) 112:112. doi:10.1016/j.cct.2021.106626
- Honea KE, Wilson KS, Fisher KL, Rubin DA. Parental and familial factors related to participation in a home-based physical activity intervention in children with obesity or Prader-Willi syndrome. *Obes Pillars* (2023) 8:8. doi:10.1016/j.obpill.2023.100084
- Putter KC, Jackson B, Thornton AL, Willis CE, Goh KMB, Beauchamp MR, et al. Perceptions of a family-based lifestyle intervention for children with overweight and

- obesity: a qualitative study on sustainability, self-regulation, and program optimization. *BMC Public Health* (2022) 22(1):1534. doi:10.1186/s12889-022-13956-6
38. Sallis JF, Prochaska JJ, Taylor WC. *A Review of Correlates of Physical Activity of Children and Adolescents* (2000). Available online at: <http://www.msse.org> (Accessed August 19, 2025).
39. Davison KK, Francis LA, Birch LL. Reexamining obesogenic families: parents' obesity-related behaviors predict girls' change in BMI. *Obes Res* (2005) 13(11):1980–90. doi:10.1038/oby.2005.243
40. Hardie JH, Landale NS. Profiles of risk: maternal health, socioeconomic status, and child health. *J Marriage Fam* (2013) 75(3):651–66. doi:10.1111/jomf.12021
41. Cameron AJ, Crawford DA, Salmon J, Campbell K, McNaughton SA, Mishra GD, et al. Clustering of obesity-related risk behaviors in children and their mothers. *Ann Epidemiol* (2011) 21(2):95–102. doi:10.1016/j.annepidem.2010.11.001
42. Julius BR, O'Shea AMJ, Francis SL, Janz KF, Laroche H. Leading by example: association between mother and child objectively measured physical activity and sedentary behavior. *Pediatr Exerc Sci* (2021) 33(2):49–60. doi:10.1123/PES.2020-0058
43. Williams J, Wake M, Hesketh K, Maher E, Waters E. Health-related quality of life of overweight and Obese children. *JAMA* (2005) 293:70–6. doi:10.1001/jama.293.1.70
44. Friedlander SL, Larkin EK, Rosen CL, Palermo TM, Redline S. Decreased quality of life associated with obesity in school-aged children. *Arch Pediatr Adolesc Med* (2003) 157:1206–11. doi:10.1001/archpedi.157.12.1206
45. Milica T, Ivana DF. Life habit of adolescents and obesity. *BMJ* (2017) 102:A91.1–A91. doi:10.1136/archdischild-2017-313273.236
46. Lier LM, Breuer C, Ferrari N, Friesen D, Maisonave F, Schmidt N, et al. Individual physical activity behaviour and group composition as determinants of the effectiveness of a childhood obesity intervention program. *Obes Facts* (2021) 14(1):100–7. doi:10.1159/000512293